東女医大誌　第83巻　臨時増刊号
頁　E675　平成25年7月

Letter to the Editor

修正：福山型先天性筋ジストロフィーにおける糖代謝
［東京女子医科大学雑誌　83（Extra）：E25～E35, 2013］

[1]東京都立府中療育センター小児科
[2]東京女子医科大学小児科
[3]東京都児童相談センター
[4]東京女子医科大学遺伝子医療センター

小峯（コミネ）　聡（サトシ）[1,2]・小平かやの（コダイラ）[3]・池谷紀代子（イケヤキヨコ）[2]・斎藤加代子（サイトウカヨコ）[2,4]・大澤真木子（オオサワマキコ）[2]

我々はこの論文に4ヵ所の修正を行った．この論文で，過去10年間に感染症で当科に入院した筋ジストロフィー40名（延べ入院回数：117回）のうち，入院時のデータを検討できた21名（FCMD：17名，DMD：4名）の年齢・血糖・CK・BMIの関係を検討した．しかし，21名の内訳は（FCMD：17名，DMD：4名）でなく，（FCMD：16名，DMD：5名）が正しく，その訂正ヵ所は26頁左段13行（目的）～14行目と26頁右段26行目（対象および方法；2. 入院筋ジストロフィー患者における検討）の2ヵ所であった．次に，対象および方法；1. 低血糖を呈したFCMDにおける検討の中で26頁左段20～21行目，「低血糖は示さなかったが尿ケトン陽性を呈したFCMD2名（症例5・症例6）（10歳10ヵ月，22歳11ヵ月）で」は，（症例5・症例6, 10歳10ヵ月，22歳11ヵ月）が正しく，括弧：）（→カンマ：，と訂正した．最後に，30頁のTable 2　The results of tolerance tests in Fukuyama congenital muscular dystrophy patients with or without hypoglycemiaの中で，Case 6のAlanine tolerance testでNormal responseをNo responseと訂正した．

Table 2　The results of tolerance tests in Fukuyama congenital muscular dystrophy patients with or without hypoglycemia

Case	Glucagon tolerance test	Alanine tolerance test	Oral glucose tolerance test
Hypoglycemia group			
1	No response	Normal response	Normal response
2	Normal response	No response	Normal response
3	Normal response	No response	Normal response
4	Normal response	No response	Normal response
Non hypoglycemia group			
5	Normal response	No response	Normal response
6	No response	No response	Normal response

Satoshi KOMINE[1,2], **Kayano KODAIRA**[3], **Kiyoko IKEYA**[2], **Kayoko SAITO**[2,4] and **Makiko OSAWA**[2]〔[1]Department of Pediatrics, Tokyo Metropolitan Fuchu Medical Center for the Disabled, [2]Department of Pediatrics, Tokyo Women's Medical University School of Medicine, [3]Tokyo Child Guidance Office, [4]Institute of Medical Genetics, Tokyo Women's Medical University〕: Erratum to: Glucose Metabolism in Patients with Fukuyama Congenital Muscular Dystrophy