

Association of Total and Differential White Blood Cell Counts with Physical Energy Expenditure

メタデータ	言語: eng 出版者: 公開日: 2018-07-17 キーワード (Ja): キーワード (En): 作成者: OYA, Junko, NAKAGAMI, Tomoko, NAITO, Yoshihiko, ENDO, Yasuhiro, UCHIGATA, Yasuko メールアドレス: 所属:
URL	http://hdl.handle.net/10470/00031905

Association of Total and Differential White Blood Cell Counts with Physical Energy Expenditure

Junko OYA¹, Tomoko NAKAGAMI¹, Yoshihiko NAITO²,
Yasuhiro ENDO³ and Yasuko UCHIGATA¹

¹Diabetes Center, Tokyo Women's Medical University School of Medicine

²School of Human Environmental Sciences, Mukogawa Women's University

³Department of Health and Community Medicine, Saitama-ken Saiseikai Kurihashi Hospital

(Accepted February 8, 2017)

Objective: To assess the association between physical energy expenditure and the total and differential white blood cell (WBC) counts which reflect systemic inflammation.

Methods: This study included 953 and 277 apparently healthy middle-aged men and women who were employed and had participated in a general health examination in 2006-2007. They completed a questionnaire regarding their usual patterns of activity during the previous month. Such behaviors included occupational, locomotive, household, leisure, and exercise activities, as well as sleep and sedentary time.

Results: In men, the total WBC count decreased with increasing levels of exercise energy expenditure, after adjusting for confounders. A similar result was observed for neutrophil counts in both genders. A multivariate logistic regression model showed that exercise energy expenditure had significant negative linear relationships with the elevated WBC and neutrophil counts in men ($p = 0.019$ and 0.026 , respectively). As compared to men who didn't exercise regularly the odds ratios of the elevated total WBC and neutrophil counts decreased significantly in the third tertile by 50 % and 51 %, respectively. However, no similar association was observed in women.

Conclusion: Exercise energy expenditure is inversely and independently associated with the WBC and neutrophil counts among healthy Japanese male workers.

Key Words: white blood cell count, physical activity, energy expenditure

Introduction

Low levels of physical activity are associated with increased risk of coronary heart disease (CHD)¹⁾, type 2 diabetes mellitus (DM)²⁾, and cardiovascular disease (CVD)-caused mortality³⁾, although the precise mechanisms remain unclear. Recently, evidence has shown that chronic inflammation is an important factor associated with the progression of

atherosclerosis⁴⁾ and insulin resistance⁵⁾. Low-grade systemic chronic inflammation is reflected by increased levels of markers such as white blood cell (WBC) count, C-reactive protein (CRP), fibrinogen, and some cytokines. WBC count is an inexpensive to measure, widely available, simple, and well-standardized biological marker of acute infection, tissue damage, and other systemic inflammatory

✉ : Tomoko NAKAGAMI Diabetes Center, Tokyo Women's Medical University, 8-1 Kawada-cho, Shinjuku-ku, Tokyo 162-8666, Japan

E-mail: nakagami.dmc@twmu.ac.jp

doi: 10.24488/jtwmu.87.Extra2_E207

Copyright © 2017 Society of Tokyo Women's Medical University

conditions. Several studies have shown that increased total WBC count is associated with the risk of developing type 2 DM⁶⁾, CHD, and all-cause mortality^{7,8)}. Regarding WBC subtypes, recent studies suggest that elevated neutrophil or granulocyte counts may be the strongest predictor of carotid arteriosclerosis, CHD, and CVD-caused mortality^{9)~11)}. Thus, the total and differential WBC counts are powerful predictors of several atherosclerotic diseases.

Previous studies have reported that higher levels of physical activity are associated with lower levels of inflammatory markers^{12)~14)}. These studies used questionnaires regarding leisure-time physical activities. However, the Japan National Health and Nutrition Survey conducted by the Japanese Ministry of Health, Labour and Welfare¹⁵⁾ reported that the 25-30 % of the Japanese population (including both genders) exercise regularly, while < 60 % attempt physical activity during their leisure time. The association between WBC count and habitual energy expenditure due to occupational, household, exercise, and leisure activities, as well as sleep and sedentary time, is unknown. Moreover, little is known about the association between differential WBC counts and physical activity levels¹⁶⁾, especially in Japanese. Thus, the purpose of this study was to assess the cross-sectional relationships between the total and differential WBC counts and daily physical activities in Japanese men and women.

Materials and Methods

Study subjects

From October 2006 to September 2007, Saitama-ken Saiseikai Kurihashi (SSK) Hospital conducted a health check-up program, in which 1,481 workers aged 25-69 years had participated, and they were followed up in the Kurihashi Lifestyle Cohort Study. Subjects were excluded if they had known DM, CVD, cancer, asthma, or certain infectious diseases, or if they were under pharmacological treatment for hypertension or dyslipidemia. Moreover, 251 subjects with abnormal WBC counts ($\geq 10,000/\mu\text{L}$ or $< 4,000/\mu\text{L}$), indicative of an infectious disease or the pathological state of leucopenia, were later ex-

cluded. The remaining 1,230 subjects (953 men and 277 women) were considered for this study.

Measurement of variables

The general health check-up procedure at SSK Hospital included biochemical laboratory tests and a self-administered questionnaire regarding smoking status, medical history, alcohol habits, and in women, menopausal status. Smoking status was classified into 3 categories (never smoked, past smoker, and current smoker). Alcohol habits were divided into 3 categories (never, occasionally, and regularly). The height, weight, blood pressure, and waist circumference (Wc) of all subjects were measured. Body height was measured to the nearest 0.1 cm with the subject standing without wearing shoes. The subjects were requested to wear light indoor clothes and the body weight was measured to the nearest 0.1 kg. Blood pressure was measured in a sitting position after 5 min of rest by using an automatic sphygmomanometer. Wc was measured to the nearest 0.1 cm at the level of the navel at the end of the expiration of a normal breath, with the subject in a standing position. Body mass index (BMI) was calculated as the body weight (in kg) divided by the body height squared (in meters). Blood samples were collected in the morning after a 10-h-long fast. The levels of fasting plasma glucose (glucose oxidase method), triglycerides (enzymatic method), and high-density lipoprotein cholesterol (direct method), and WBC counts (Kinetic WBC optical count, CELL-DYN; Abbott Japan, Tokyo) were measured at the hospital laboratory. Insulin concentrations were measured by an immunoradiometric assay at a commercial laboratory.

Physical activities

The participants completed the Japan Arteriosclerosis Longitudinal Study Physical Activity Questionnaire (JALSPAQ)^{17,18)} regarding their usual patterns of physical activity during the previous month. The questionnaire comprises 14 questions on sleep time, occupation, locomotion, housework, leisure-time physical activities (e.g., gardening, home carpentry, car washing), and exercise. Questionnaire data were converted using the intensity of each physical activity expressed in metabolic

equivalents (METs), according to the Compendium by Ainsworth et al¹⁹⁾, and summarized as METs · h/day and energy expenditure. In this questionnaire, occupational physical activity was assessed by recording the duration of sitting (1.5 METs), standing (2.0 METs), and walking (3.0 METs). Similarly, locomotional physical activity was assessed by walking (3.0 METs) and cycling (4.0 METs); housework physical activity was assessed by cooking (2.3 METs), washing (2.0 METs), cleaning (3.5 METs), and nursing or providing child care (3.0 METs). Leisure-time and exercise physical activities were calculated by each selected physical activity value (METs) multiplied by activity duration. The JALSPAQ values were validated against the “gold standard” doubly labeled water (DLW) method by using the data from 226 Japanese men and women aged 20-83 years²⁰⁾. JALSPAQ slightly underestimated total energy expenditure: the difference in the mean and standard error was $-1.15 \pm 1.92 \text{ kcal} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$. The total energy expenditure values obtained using JALSPAQ and DLW were moderately correlated (Spearman's correlation coefficient = 0.742, $p < 0.001$; intraclass correlation coefficient = 0.648, $p < 0.001$), and the 95 % limit of agreement was 4.99 to 2.69 kcal/kg. The energy expenditure of each physical activity ($\text{kcal} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$) in JALSPAQ was validated by the 24-hour weighed physical activity record among 122 Japanese participants (average age, 55.8 years). Spearman's correlation coefficients were as follows: 0.75 for occupational physical activity, 0.13 for locomotional physical activity, 0.59 for housework physical activity, 0.60 for exercise physical activity, and 0.41 for leisure-time physical activity¹⁷⁾.

Statistical analysis

Statistical analyses were performed and reported separately for men and women. The chi-square test was used to compare proportions and the student's t-test was used to compare the mean values between genders. Because of the skewed distribution of triglyceride, we used log-transformed values in the analyses and showed the medians and interquartile ranges. The Mann-Whitney's U test was used to compare the values of each energy expendi-

ture. Spearman's correlation coefficients were used to examine the correlation between the total and differential WBC counts and each of the daily energy expenditures. Participants were categorized into 4 groups of energy expenditure in selected physical activities that showed significant correlation between the total and differential WBC counts (total energy expenditure in both genders [Quartile: < 31.3 , $31.4-33.0$, $33.1-36.7$, $\geq 36.8 \text{ kcal} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ in men; < 32.3 , $32.4-33.8$, $33.9-37.2$, $\geq 37.3 \text{ kcal} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ in women] and exercise in both genders [0 and tertile: 0, < 0.69 , $0.70-1.48$, $\geq 1.49 \text{ kcal} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ in men; 0, < 0.54 , $0.55-1.38$, $\geq 1.39 \text{ kcal} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ in women]). Sleep energy expenditure was not selected given its discrete distribution related to questionnaire framing. The general linear model was used to test the equality of means of total WBC and differential counts across the 4 groups of energy expenditure. The crude and multivariate adjusted odds ratios (ORs) and their 95 % confidence intervals (CIs) associated with the highest quartile of total WBC and neutrophil counts were calculated for each category of total or selected physical activity by using logistic regression models. In the multivariate model, age (continuous), BMI (continuous), Wc (continuous), alcohol habits (categorical), smoking status (categorical), fasting insulin concentrations, and menopause status (in women only; categorical) were included as confounding factors. To confirm the causal impact of physical activity levels on WBC counts in this cross-sectional study, the equality of means of physical activities across quartiles of WBC counts were examined.

The Statistical Package for Social Sciences (SPSS) for Windows (version 21.0, Chicago, IL, USA) was used for all statistical analyses. All reported p values are two-tailed, and $p < 0.05$ was considered statistically significant. The study was approved by the Institutional Review Board of SSK Hospital, and informed consent was obtained from the study participants.

Results

Characteristics of study subjects

Overall, the male subjects were more obese, had worse CVD risk profiles, and had a higher preva-

Table 1 Characteristics of study subjects

	Men (n = 953)	Women (n = 277)	p value
Age (years)	49 ± 8	49 ± 8	0.837
Body mass index (kg/m ²)	23.7 ± 3.2	22.8 ± 3.2	< 0.001
Waist circumference (cm)	85.4 ± 8.3	81.5 ± 8.6	< 0.001
Systolic blood pressure (mmHg)	124 ± 15	120 ± 16	< 0.001
Diastolic blood pressure (mmHg)	80 ± 12	72 ± 12	< 0.001
High-density lipoprotein cholesterol (mmol/L)	1.40 ± 0.36	1.68 ± 0.39	< 0.001
Triglyceride (mmol/L)	1.16 (0.87, 1.60)	0.89 (0.64, 1.22)	< 0.001
Fasting plasma glucose (mmol/L)	5.3 ± 0.7	5.1 ± 0.8	< 0.001
Fasting insulin (μU/mL)	6.9 ± 5.9	6.1 ± 3.4	0.008
Alcohol intake regularly (%)	40.6	11.1	0.799
Current smoker (%)	65.6	15.9	< 0.001
Menopause (%)	—	49.2	
Total white blood cell count (/μL)	5,798 ± 1,203	5,320 ± 1,013	< 0.001
Energy expenditure (kcal·kg ⁻¹ ·day ⁻¹)			
Total	33.6 (31.5, 37.3)	35.2 (32.6, 38.6)	< 0.001
During Sleep	6.8 (6.0, 7.1)	6.4 (6.0, 7.1)	< 0.001
Occupation	10.7 (8.6, 16.4)	8.5 (5.0, 13.5)	< 0.001
Locomotion	1.5 (0.5, 3.0)	1.2 (0.5, 2.5)	< 0.001
Household	0.2 (0.0, 0.8)	4.9 (3.0, 6.8)	0.032
Exercise	0.3 (0.0, 1.2)	0.1 (0.0, 0.8)	< 0.001
Leisure-time	0.2 (0.0, 0.7)	0.2 (0.0, 0.9)	0.006
Sedentary	11.6 (9.1, 13.6)	12.1 (9.5, 14.6)	0.223

Variables are the mean ± standard deviation or median and interquartile range for continuous variables and percentages of subjects for categorical variables.

Table 2 Spearman's correlation coefficients (γ) between total white blood cell count and each type of energy expenditure

	Men		Women	
	γ	p value	γ	p value
Total WBC count				
Energy expenditure (kcal·kg ⁻¹ ·day ⁻¹)				
Total	0.010	0.766	-0.060	0.323
During sleep	-0.071	0.029	0.021	0.730
Occupational	0.028	0.385	0.035	0.567
Locomotion	-0.044	0.174	0.020	0.741
Household	0.048	0.136	-0.077	0.200
Exercise	-0.072	0.026	-0.159	0.008
Leisure-time	-0.006	0.846	-0.002	0.972
Sedentary	0.008	0.809	-0.050	0.404
Neutrophil count				
Energy expenditure (kcal·kg ⁻¹ ·day ⁻¹)				
Total	-0.028	0.496	-0.028	0.704
During sleep	-0.068	0.091	-0.030	0.684
Occupational	0.035	0.392	0.084	0.260
Locomotion	-0.018	0.096	-0.006	0.939
Household	-0.048	0.238	-0.133	0.073
Exercise	-0.085	0.035	-0.007	0.928
Leisure-time	-0.001	0.759	0.040	0.593
Sedentary	0.012	0.759	-0.055	0.465

WBC, white blood cell.

lence of current smokers compared to the female subjects. With respect to energy expenditure, men, when compared to women, were physically more

active in occupational, locomotion, and exercise activities, while physically less active in household activities and in all activities taken together (Table 1).

Table 3 Adjusted means of total and differential white blood cell counts in the four groups of each of physical energy expenditure

	Group 1	Group 2	Group 3	Group 4	p for linear trend
Men					
Total	240	240	237	236	
Energy expenditure (kcal·kg ⁻¹ ·day ⁻¹)	<31.3	31.4–33.0	33.1–36.7	≥36.8	
Total WBC (/μL)	5,857 ± 1,187	5,807 ± 1,252	5,738 ± 1,160	5,792 ± 1,216	0.759
Exercise					
Number	428	172	177	176	
Energy expenditure (kcal·kg ⁻¹ ·day ⁻¹)	0	<0.69	0.70–1.48	≥1.49	
Total WBC (/μL)	5,932 ± 1,257	5,748 ± 1,117	5,653 ± 1,111 *	5,628 ± 1,088 *	0.012
Neutrophil (/μL)	3,430 ± 905	3,256 ± 790	3,251 ± 810	3,176 ± 630 *	0.033
Lymphocyte (/μL)	1,974 ± 604	1,961 ± 626	1,902 ± 550	1,819 ± 515	0.122
Monocyte (/μL)	341 ± 105	318 ± 90	326 ± 88	316 ± 91	0.061
Eosinophil (/μL)	207 ± 141	207 ± 122	190 ± 145	206 ± 131	0.648
Basophil (/μL)	31 ± 19	32 ± 20	28 ± 17	28 ± 15	0.241
Women					
Total	70	70	68	69	
Energy expenditure (kcal·kg ⁻¹ ·day ⁻¹)	<32.3	32.4–33.8	33.9–37.2	≥37.3	
Total WBC (/μL)	5,383 ± 1,036	5,441 ± 1,183	5,213 ± 792	5,241 ± 1,000	0.487
Exercise					
Number	157	40	41	39	
Energy expenditure (kcal·kg ⁻¹ ·day ⁻¹)	0	<0.54	0.55–1.38	≥1.39	
Total WBC (/μL)	5,290 ± 1,094	5,520 ± 1,141	5,239 ± 770	4,968 ± 627	0.253
Neutrophil (/μL)	2,957 ± 722	3,405 ± 818	2,941 ± 728	2,667 ± 575	0.032
Lymphocyte (/μL)	1,793 ± 522	1,624 ± 506	1,737 ± 410	1,676 ± 340	0.342
Monocyte (/μL)	271 ± 90	287 ± 93	247 ± 118	223 ± 120	0.066
Eosinophil (/μL)	179 ± 133	175 ± 138	125 ± 115	158 ± 154	0.135
Basophil (/μL)	28 ± 20	31 ± 18	22 ± 18	27 ± 30	0.261

WBC, white blood cell. Variables are the mean ± standard error.

Adjusted for age, body mass index, waist circumference, smoking status, alcohol habits, fasting insulin concentrations in both men and women, and only in women, menopausal status.

*p<0.05 vs group 1.

Four groups: Total energy expenditure in both genders [Quartile: <31.3, 31.4–33.0, 33.1–36.7, ≥36.8, kcal·kg⁻¹·day⁻¹ in men; <32.3, 32.4–33.8, 3.9–37.2, ≥37.3 kcal·kg⁻¹·day⁻¹ in women] and exercise in both genders [0 and tertile: 0, <0.69, 0.70–1.48, ≥1.49 kcal·kg⁻¹·day⁻¹ in men; 0, <0.54, 0.55–1.38, ≥1.39 kcal·kg⁻¹·day⁻¹ in women].

Correlation between energy expenditure and WBC count

Table 2 shows that energy expenditure during sleep time negatively correlated with the total WBC count in men, and exercise energy expenditure negatively correlated with the total WBC counts in both genders. The exercise energy expenditure was inversely correlated with the neutrophil count in men. No significant correlation was found between other differential WBC counts and each form of energy expenditure in either gender.

Total and differential WBC counts and physical energy expenditure

No significant relationships were found in the total WBC counts across quartiles of total energy expenditure in either gender. There were also no significant associations between the total WBC counts and the 4 groups of energy expenditure during sleep time. The total WBC count decreased with increasing levels of exercise energy expenditure, after adjusting for confounding factors in men. Similar results appeared for neutrophil counts in both genders (Table 3).

The multivariate logistic regression model

Table 4 Odds ratios and their 95% confidence intervals for the total and exercise energy expenditure associated with the highest quartile of total WBC and neutrophil counts

	Total energy expenditure				p for linear trend
	1st Quartile	2nd Quartile	3rd Quartile	4th Quartile	
Men					
Number	240	240	237	236	
Energy expenditure (kcal·kg ⁻¹ ·day ⁻¹)	<31.3	31.4–33.0	33.1–36.7	≥36.8	
WBC					
Crude OR (95%CI)	1.00 (ref)	0.82 (0.55–1.25)	0.86 (0.57–1.29)	0.89 (0.60–1.35)	0.234
Multivariate OR (95%CI)	1.00 (ref)	0.75 (0.48–1.17)	0.88 (0.57–1.36)	0.82 (0.53–1.28)	0.634
Women					
Number	70	70	68	69	
Energy expenditure (kcal·kg ⁻¹ ·day ⁻¹)	<32.3	32.4–33.8	33.9–37.2	≥37.3	
WBC					
Crude OR (95%CI)	1.00 (ref)	2.07 (0.93–4.63)	1.74 (0.76–3.96)	1.71 (0.75–3.88)	0.247
Multivariate OR (95%CI)	1.00 (ref)	3.10 (0.93–10.29)	2.02 (0.65–6.30)	2.94 (0.91–9.54)	0.910
	Exercise energy expenditure				p for linear trend
	Group 1	Group 2	Group 3	Group 4	
Men					
Number	428	172	177	176	
Energy expenditure (kcal·kg ⁻¹ ·day ⁻¹)	0	<0.69	0.70–1.48	≥1.49	
WBC					
Crude OR (95%CI)	1.00 (ref)	0.75 (0.50–1.11)	0.60 (0.39–0.91)	0.48 (0.30–0.75)	0.004
Multivariate OR (95%CI)	1.00 (ref)	0.87 (0.53–1.41)	0.68 (0.42–1.12)	0.50 (0.27–0.91)	0.019
Neutrophil					
Crude OR (95%CI)	1.00 (ref)	0.75 (0.45–1.25)	0.66 (0.35–1.22)	0.42 (0.19–0.93)	0.015
Multivariate OR (95%CI)	1.00 (ref)	0.87 (0.48–1.55)	0.68 (0.31–1.22)	0.49 (0.26–0.82)	0.026
Women					
Number	157	40	41	39	
Energy expenditure (kcal·kg ⁻¹ ·day ⁻¹)	0	<0.54	0.55–1.38	≥1.39	
WBC					
Crude OR (95%CI)	1.00 (ref)	1.01 (0.46–2.19)	0.97 (0.45–2.11)	0.30 (0.10–0.90)	0.193
Multivariate OR (95%CI)	1.00 (ref)	1.68 (0.63–4.43)	1.27 (0.45–3.61)	0.34 (0.07–1.66)	0.318
Neutrophil					
Crude OR (95%CI)	1.00 (ref)	2.44 (0.66–9.08)	0.98 (0.30–3.26)	0.61 (0.12–3.15)	0.115
Multivariate OR (95%CI)	1.00 (ref)	1.22 (0.40–3.75)	0.98 (0.33–2.91)	0.62 (0.20–1.96)	0.506

WBC, white blood cell; OR, odds ratio; CI, confidence interval.

Adjusted for age, body mass index, waist circumference, smoking status, alcohol habits, fasting insulin concentrations in both men and women, and menopausal status only in women.

Four groups: Total energy expenditure [Quartile: <31.3, 31.4–33.0, 33.1–36.7, ≥36.8 kcal·kg⁻¹·day⁻¹ in men; <32.3, 32.4–33.8, 33.9–37.2, ≥37.3 kcal·kg⁻¹·day⁻¹ in women] and exercise [0 and tertile: 0, <0.69, 0.70–1.48, ≥1.49 kcal·kg⁻¹·day⁻¹ in men; 0, <0.54, 0.55–1.38, ≥1.39 kcal·kg⁻¹·day⁻¹ in women].

showed that exercise energy expenditure had statistically significant linear negative relationships with the WBC and neutrophil counts in men ($p = 0.019$ and 0.026 , respectively, for the linear trends)

(Table 4). The adjusted OR related to the highest quartiles of the total WBC and neutrophil counts decreased significantly in group 4 (3rd tertile) of exercise energy expenditure (≥ 1.49 kcal · kg⁻¹ ·

day⁻¹) when compared to group 1 (no habitual exercise) of exercise energy expenditure by 50 % and 51 %, respectively (Table 4). In contrast, no significant association was found between exercise energy expenditure and the highest quartiles of the WBC and neutrophil counts in women.

In contrast, the WBC counts did not predict the physical activity levels in either gender (data not shown).

Discussion

The current study has shown that the total WBC and neutrophil counts, which reflect chronic inflammatory conditions, decreased with increasing levels of exercise energy expenditure in men, after adjusting for confounding factors. In women, only the neutrophil count decreased with increasing levels of exercise energy expenditure. In Japan, few people exercise regularly¹⁵⁾, and physical activity (including non-exercise activity) thermogenesis was examined in this study. However, no association was found between daily physical activity excluding exercise and total WBC and its differential counts. Thus, our results indicated a dose-responsive negative association between exercise and total WBC and neutrophil counts among healthy Japanese male workers.

Previous cross-sectional studies have shown an association between the total WBC count and physical activity. Geffken et al¹²⁾ reported that persons in the highest quartile of leisure-time physical activity ($\geq 2,270$ kcal/week) had 6 % lower concentrations of total WBCs compared with those in the lowest quartile (< 367.5 kcal/week) in a healthy elderly population. Wannamethee et al¹³⁾ showed a significant and inverse dose-response relationship between physical activity (walking, cycling, recreational activity, and sports) and total WBC count in elderly men. Abramson and Vaccarino¹⁴⁾ have reported that a higher frequency of physical activity was independently associated with lower ORs for elevated total WBC count among healthy adults. These reports were consistent with our findings.

Recent studies have shown that elevated neutrophil counts may be the strongest predictor of CVD, CHD, and death^{11/21)}. Johannsen et al¹⁶⁾ reported that

higher levels of cardiorespiratory fitness were associated with lower counts of total WBC, neutrophils, lymphocytes, and basophils. In contrast, another study that assessed the association between maximal oxygen uptake and the subtypes of WBC showed that monocyte count, not neutrophil count, was related to aerobic capacity²²⁾. Our results show that higher levels of exercise energy expenditure may decrease the neutrophil count, leading to reduced risk of CVD.

Despite evidence for an association between the total and differential WBC counts and physical activity, the underlying biological mechanisms remain unclear. Proinflammatory cytokines such as tumor necrosis factor- α (TNF- α) and interleukin (IL)-6, which are secreted from the adipose tissue, are known to increase WBC counts²³⁾. TNF- α is a potent stimulator of IL-6 production, and IL-6 directly stimulates neutrophils²⁴⁾. Several studies have demonstrated that habitual exercise was inversely related to both TNF- α and IL-6 levels^{25/26)}. In animal model, it has shown that IL-6 derived from skeletal muscle by acute exercise mediate the increase of leptin and insulin sensitivity in hypothalamus²⁷⁾. These findings indicated that exercise could have appetite-suppressive actions via hypothalamus and might lead decreasing adipocyte which was associated with chronic inflammation. Moreover, exercise intervention reduced the levels of IL-8, a neutrophil chemokine derived from the adipose tissue²⁸⁾, which subsequently contributed to the migration of granulocytes and decreased the levels of granulocyte colony-stimulating factor, monocyte chemoattractant protein-1 (MCP-1)²⁹⁾. Recently, it has known that one of the effects of exercise on muscle is mediated by the transcriptional coactivator peroxisome proliferator-activated receptor γ coactivator-1 α (PGC-1 α). PGC-1 α expression in muscle stimulates secretion of new hormone, irisin. It is shown that irisin drives browning of white fat and thermogenesis³⁰⁾. These changes of adipose tissue may reduce circulating leptin which has been shown to activate neutrophils³¹⁾. Thus, exercise may be important to suppress chronic systemic inflammation, which leads to CVD, among Japanese male workers, who

rarely exercise regularly. It has been examined that exercise influenced the immunological parameters through alteration of autonomic nervous system, however, the association between WBC counts and autonomic nerve activity is unknown.

There is a gender-based difference in the progression of DM and CVD³²⁾. A similar gender bias has been reported in the total and differential WBC counts³³⁾. In the current study, we were unable to find a significant association between exercise energy expenditure and the elevated WBC and neutrophil counts in women. The reasons for this are unclear. The number of women in our study was less than one-third the number of men, thus our findings in women might be weakened by the smaller sample size. There may also be environmental gender-based differences in lifestyle such as smoking status. In our study, the male subjects included a higher proportion of smokers than the female subjects. In addition, a female sex hormone, estrogen, may protect against atherosclerosis by decreasing inflammatory cell adhesion³⁴⁾. The menstrual cycle also affects WBC counts. It is known that granulocyte counts are higher in the luteal phase compared to the follicular phase³⁵⁾. Gender-based differences are also noted in the subcutaneous and visceral fat distribution. It was reported that visceral adipose tissue produced more MCP-1 than did subcutaneous adipose tissue³⁶⁾. Therefore, visceral fat is a more important inducer of low-grade inflammation than subcutaneous fat. As women are known to have more subcutaneous fat and less visceral fat than men, these differences might influence our results.

There were several limitations in our study. First, this study had a cross-sectional design, precluding the establishment of temporal relationships between the total and differential WBC counts and activity energy expenditure. Further prospective data analysis is needed to confirm our findings. Second, the results of the questionnaire-based survey for alcohol consumption and smoking status were self-reported and not validated. Third, the JALSPAQ has been validated in prior studies¹⁷⁾⁽²⁰⁾; however, participants of these validation studies were

limited to middle-aged adults, and the degree of correlation between estimated energy expenditure in JALSPAQ and DLW or 24-hour record was moderate. Validation studies utilizing a combination of accelerometry and 24-hour record study spanning various ages are necessary for accurate estimation of physical activity level. Fourth, high-sensitivity C-reactive protein (hs-CRP) is known as a sensitive marker for the inflammatory component of metabolic syndrome and CVD³⁷⁾⁽³⁸⁾. However, acquiring CRP data was cost-prohibitive, whereas WBC count is a stable, well-standardized, and inexpensive marker of routine health care data. Further research is needed to examine the relationships between a more sensitive inflammatory marker and physical activity. In addition to its limitations, our study has some major advantages. First, we excluded subjects whose WBC counts exceeded 10,000/ μ L or were $<4,000/\mu$ L, limiting our subjects to individuals whose WBC counts were within the normal range. Second, to avoid systematic bias, we were extremely careful in data collection. We excluded subjects who had a medical history of diseases associated with low-grade inflammation.

Conclusion

The exercise energy expenditure is inversely and independently associated with the WBC and neutrophil counts among healthy Japanese male workers. Further prospective data analysis is needed to confirm our findings.

Acknowledgments

This study was supported by grants from the Japanese Ministry of Health, Labour, and Welfare; the Japan Medical Women's Association; the Tokyo Women's Medical University Association; the Yayoi Yoshioka Research Fund; the Yazuya Food and Health Research Foundation; the Japan Diabetes Society; and the Japan Automobile Manufacturers Association, Inc. for T.N.

COMPETING INTERESTS: None declared.

References

- 1) Folsom AR, Arnett DK, Hutchinson RG et al: Physical activity and incidence of coronary heart disease in middle-aged women and men. *Med Sci*

- Sports Exerc **29**: 901–909, 1997
- 2) **Manson JE, Rimm EB, Stampfer MJ et al**: Physical activity and non-insulin-dependent diabetes mellitus in women. *Lancet* **338**: 774–778, 1991
 - 3) **Smith GD, Shipley MJ, Batty GD et al**: Physical activity and cause-specific mortality in the Whitehall study. *Public Health* **114**: 308–315, 2000
 - 4) **Libby P**: Inflammation in atherosclerosis. *Nature* **420**: 868–874, 2002
 - 5) **De Rooij SR, Nijpels G, Nilsson PM et al**: Low-grade chronic inflammation in the relationship between insulin sensitivity and cardiovascular disease (RISC) population: associations with insulin resistance and cardiometabolic risk profile. *Diabetes Care* **32**: 1295–1301, 2009
 - 6) **Nakanishi N, Yoshida H, Matsuo Y et al**: White blood-cell count and the risk of impaired fasting glucose or type II diabetes in middle-aged Japanese men. *Diabetologia* **45**: 42–48, 2002
 - 7) **Weijenberg MP, Feskens EJ, Kromhout D**: White blood cell count and the risk of coronary heart disease and all-cause mortality in elderly men. *Arterioscler Thromb Vasc Biol* **16**: 499–503, 1996
 - 8) **Leng SX, Xue QL, Huang Y et al**: Baseline total and differential white blood cell counts and 5-year all-cause mortality in community-dwelling older women. *Exp Gerontol* **40**: 982–987, 2005
 - 9) **Huang ZS, Jeng JS, Wang CH et al**: Correlations between peripheral differential leukocyte counts and carotid atherosclerosis in non-smokers. *Atherosclerosis* **158**: 431–436, 2001
 - 10) **Rana JS, Boekholdt SM, Ridker PM et al**: Differential leucocyte count and the risk of future coronary artery disease in healthy men and women: the EPIC-Norfolk prospective population study. *J Intern Med* **262**: 678–689, 2007
 - 11) **Gillum RF, Mussolino ME, Madans JH**: Counts of neutrophils, lymphocytes, and monocytes, cause-specific mortality and coronary heart disease: the NHANES-I epidemiologic follow-up study. *Ann Epidemiol* **15**: 266–271, 2005
 - 12) **Geffken DF, Cushman M, Burke GL et al**: Association between physical activity and markers of inflammation in a healthy elderly population. *Am J Epidemiol* **153**: 242–250, 2001
 - 13) **Wannamethee SG, Lowe GD, Whincup PH et al**: Physical activity and hemostatic and inflammatory variables in elderly men. *Circulation* **105**: 1785–1790, 2002
 - 14) **Abramson JL, Vaccarino V**: Relationship between physical activity and inflammation among apparently healthy middle-aged and older US adults. *Arch Intern Med* **162**: 1286–1292, 2002
 - 15) **The Japanese Ministry of Health, Labor and Welfare**: Section 2: Settings of anthropometry and lifestyle. In *The State of National Health and Nutrition- Report from Examination of National Health and Nutrition*, pp224–225, Dai-ichi shuppan, Tokyo (2009) (in Japanese)
 - 16) **Johannsen NM, Priest EL, Dixit VD et al**: Association of white blood cell subfraction concentration with fitness and fatness. *Br J Sports Med* **44**: 588–593, 2010
 - 17) **Naito Y, Harada A, Inoue S et al**: Report of the physical activity research of the Japan Arteriosclerosis Longitudinal Study. *Research in Exercise Epidemiology* **5**: 1–7, 2003 (in Japanese)
 - 18) **Japan Arteriosclerosis Longitudinal Study (JALS) Group**: Japan Arteriosclerosis Longitudinal Study-Existing Cohorts Combine (JALS-ECC): rationale, design, and population characteristics. *Circ J* **72**: 1563–1568, 2008
 - 19) **Ainsworth BE, Haskell WL, Whitt MC et al**: Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc* **32**: S498–S516, 2000
 - 20) **Ishikawa-Takata K, Naito Y, Tanaka S et al**: Use of doubly labeled water to validate a physical activity questionnaire developed for the Japanese population. *J Epidemiol* **21**: 114–121, 2011
 - 21) **Horne BD, Anderson JL, John JM et al**: Which white blood cell subtypes predict increased cardiovascular risk? *J Am Coll Cardiol* **45**: 1638–1643, 2005
 - 22) **Michishita R, Shono N, Inoue T et al**: Associations of monocytes, neutrophil count, and C-reactive protein with maximal oxygen uptake in overweight women. *J Cardiol* **52**: 247–253, 2008
 - 23) **Sim E**: Humoral factors (Natural immune system). Oxford Univ Press, Oxford (1993)
 - 24) **McCarty MF**: Interleukin-6 as a central mediator of cardiovascular risk associated with chronic inflammation, smoking, diabetes, and visceral obesity: down-regulation with essential fatty acids, ethanol and pentoxifylline. *Med Hypotheses* **52**: 465–477, 1999
 - 25) **Tsukui S, Kanda T, Nishino M et al**: Moderate-intensity regular exercise decreases serum tumor necrosis factor-alpha and HbA1c levels in healthy women. *Int J Obes Relat Metab Disord* **24**: 1207–1211, 2000
 - 26) **Smith JK, Dykes R, Douglas JE et al**: Long-term exercise and atherogenic activity of blood mononuclear cells in persons at risk of developing ischemic heart disease. *JAMA* **281**: 1722–1727, 1999
 - 27) **Flores MB, Fernandes MF, Ropelle ER et al**: Retracted Article Exercise improves insulin and leptin sensitivity in hypothalamus of Wistar rats. *Diabetes* **55**: 2554–2561, 2006
 - 28) **Troscid M, Lappegård KT, Claudi T et al**: Exercise reduces plasma levels of the chemokines MCP-1 and IL-8 in subjects with metabolic syndrome. *Eur Heart J* **25**: 349–355, 2004
 - 29) **Adamopoulos S, Parissis J, Kroupis C et al**: Physical training reduces peripheral markers of inflammation in patients with chronic heart failure. *Eur Heart J* **22**: 791–797, 2001
 - 30) **Bostrom P, Wu J, Jedrychowski MP et al**: A PGC 1 α -dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature* **481**: 463–468, 2012
 - 31) **Zarkesh-Esfahani H, Pockley AG, Wu Z et al**: Leptin indirectly activates human neutrophils via

- induction of TNF- α . J Immunol **172**: 1809–1814, 2004
- 32) **Onat A, Hergenç G, Keleş I et al**: Sex difference in development of diabetes and cardiovascular disease on the way from obesity and metabolic syndrome. Metabolism **54**: 800–808, 2005
- 33) **Bain BJ**: Ethnic and sex differences in the total and differential white blood cell count and platelet count. J Clin Pathol **49**: 664–666, 1996
- 34) **Mendelsohn ME, Karas RH**: Molecular and cellular basis of cardiovascular gender differences. Science **308**: 1583–1587, 2005
- 35) **Faas M, Bouman A, Moesa H et al**: The immune response during the luteal phase of the ovarian cycle: a TH2-type response? Fertil Steril **74**: 1008–1013, 2000
- 36) **Bruun JM, Lihn AS, Pedersen SB et al**: Monocyte chemoattractant protein-1 release is higher in visceral than subcutaneous human adipose tissue (AT): implication of macrophages resident in the AT. J Clin Endocrinol Metab **90**: 2282–2289, 2005
- 37) **Oda E, Kawai R**: Comparison between high-sensitivity C-reactive protein (hs-CRP) and white blood cell count (WBC) as an inflammatory component of metabolic syndrome in Japanese. Intern Med **49**: 117–124, 2010
- 38) **Ridker PM, Wilson PW, Grundy SM**: Should C-reactive protein be added to metabolic syndrome and to assessment of global cardiovascular risk? Circulation **109**: 2818–2825, 2004

身体消費エネルギー量と総白血球数，白血球分画の関係

¹東京女子医科大学糖尿病センター（糖尿病・代謝内科）

²武庫川女子大学生生活環境学部食物栄養学科

³埼玉県済生会栗橋病院・健診部

オオヤ ジュンコ ナカガミ トモコ ナイトウ ヨシヒコ エンドウ ヤスヒロ ウチガタ ヤスコ
大屋 純子¹・中神 朋子¹・内藤 義彦²・遠藤 康弘³・内潟 安子¹

〔背景〕総白血球数（WBCC）上昇は心血管疾患（CVD）予測の簡便な炎症マーカーである。身体活動量の増加により CVD 発症予防が期待されるが、身体活動量と WBCC とその分画の関連についての報告は少ない。

〔目的〕各種身体活動の 1 日の体重当たり消費エネルギー量（EE）と WBCC，白血球分画との関連を横断的に検討する。

〔方法〕対象は、就業中の 25-69 歳の非糖尿病、高血圧や脂質異常症の薬物療法を受けていないドック受診者 1,230 名（男性 953 名、平均年齢 49 ± 8 歳、平均 BMI 23.5 ± 3.3 kg/m²）である。身体活動質問票を用いて過去 1 か月間の身体活動内容を調査し、男女別に睡眠、仕事、運動、非活動、合計の EE を算出した。EE と WBCC の間で相関を認めた活動ごとに、WBCC か白血球分画高値（4 分割の最上位）を目的変数、活動の EE を説明変数としたロジスティック回帰モデルを用いて、活動の EE（G1-4 に 4 分割）の独立した関連性を分析した。

〔結果〕合計 EE は WBCC と一定の傾向を示さなかったが、男女ともに運動 EE と WBCC は負の相関、男性でのみ睡眠 EE と WBCC に負の相関を認めた。睡眠 EE の 4 分位で WBCC，白血球分画には一定の傾向は見られなかったが、運動 EE の 4 分割（運動なし：G1，する者で 3 分位：G2-4）では、仕事 EE，睡眠 EE，年齢，BMI，臍周囲径，飲酒，喫煙，インスリン濃度，女性ではさらに閉経で調整後も、男性では WBCC と好中球数が、女性では好中球数が、活動量の増加にともなって低下した。男性で、運動 EE の WBCC 高値に対するオッズ比は、G1 と比較し G2→G4 に向け、0.87, 0.68, 0.50 (Trend $p=0.019$)，好中球高値に対するオッズ比は 0.87, 0.68, 0.49 (Trend $p=0.026$) であった。同様の解析を女性で行ったが有意な傾向はなかった。男性では運動 EE が 1.49 kcal/kg/day (=G4) 以上で WBCC，好中球数高値のリスクが有意に上昇していた。

〔総括〕男性では運動 EE と白血球数，特に好中球数との負の関連性が認められた。