

Immunohistochemical characterization of
interstitial cells of Cajal (ICC) in the mouse
intestine using whole mount preparations with
particular reference to their density,
morphological features, and network patterns

メタデータ	言語: jpn 出版者: 公開日: 2021-02-02 キーワード (Ja): キーワード (En): 作成者: 川島, 章子 メールアドレス: 所属:
URL	https://doi.org/10.20780/00032712

Immunohistochemical characterization of interstitial cells of Cajal (ICC) in the mouse intestine using whole mount preparations – with particular reference to their density, morphological features, and network patterns

Shoko Kawashima^{1,2}, Sachiko Miyamoto-Kikuta^{2,3*}, Taichi Ezaki², and Takahiro Okamoto¹

¹Department of Breast, Endocrine, and Pediatric Surgery, ²Department of Anatomy and Developmental Biology, ³Department of Anatomy, Division of Microscopic Anatomy and Developmental Morphology, Tokyo Women's Medical University (TWMU), 8-1 Kawada-cho, Shinjuku-ku, Tokyo 162-8666, Japan

Summary. The aim of this study was to evaluate the distribution pattern of the different subtypes of the interstitial cells of Cajal (ICC), to understand their roles in gastrointestinal motility and interaction with the enteric nervous system depending on their location. Based on morphological analyses using the immunohistochemistry of whole mount preparations of mouse intestine, we investigated the normal local density and morphological features of ICC and statistically compared the number of the cells between the mesenteric and anti-mesenteric sides, and between the proximal and distal sites. Density gradients were shown in ICC associated with the deep muscular plexus, the myenteric plexus in the large intestine, and the longitudinal muscle. However, such

changes in cell density were not always proportional to other subtypes, even in the same part of the intestine. As the pacemaker cells for intestinal motility, dense and seamless network connections were shown in ICC associated with the myenteric plexus in the small intestine and ICC associated with the submuscular plexus in the large intestine without any gradient of cell density, in contrast with the difference of morphological network patterns between proximal and distal sites. Thus, we conclude that the distribution patterns of ICC differ in cell density, morphological features, and network patterns, depending on the individual subtype and site of the intestine. Therefore, careful studies on the quantification and distribution of ICC under normal conditions should be important in diagnosing clinical disorders related to abnormal distribution of ICC.

Received December 1, 2019

Address for correspondence: Sachiko Miyamoto-Kikuta

³Department of Anatomy, Division of Microscopic Anatomy and Developmental Morphology, Tokyo Women's Medical University, 8-1, Kawada-cho, Shinjuku-ku, Tokyo 162-8666, Japan

Tel & Fax: +81-3-5269-7407

E-mail: kikuta.sachiko@twmu.ac.jp

Keywords: Interstitial cells of Cajal (ICC), subtype, distribution, intestines, mouse

Introduction

Transport and absorption of food and fluid are highly regulated by constitutive gastrointestinal motility. It has been

shown that several components of the regulatory mechanism of smooth muscle movement may be involved (Sanders *et al.* 2012).

As the main control system for gastrointestinal motility, the enteric nervous system (ENS) has been studied in great detail (Costa *et al.* 1971; Furness and Costa 1971, 1980). Quantitative studies of ENS have been based on morphological observations of neurons in the myenteric plexus (Irwin 1931; Gabella 1971). These studies have revealed that the distribution of neurons in the gut is not uniform but differs in different parts. For instance, the large intestine contains a greater density of neurons than the small intestine and their mesenteric sides have a greater density than the anti-mesenteric sides. Therefore, gastrointestinal motility differs in different parts of the alimentary tract, mainly due to the uneven distribution of the ENS. For example, in the small intestine, peristaltic movement is uni-directional (Cannon 1902) and segmentation motor activity is a movement that facilitates absorption of nutrients. In human, the movement of the large intestine has been classified as non-propulsive segmentation, propulsive, and retropropulsive (antiperistalsis) movements (Ritchie 1970). In mammals, antiperistalsis is limited to the proximal part of the colon (Elliott and Barclay-Smith 1904).

Another control system, the interstitial cells of Cajal (ICC), have been shown to play an important role in gastrointestinal motility, by many studies (Huizinga *et al.* 1995; Sanders *et al.* 2014). Morphological studies using electron microscopy have elucidated the microstructure of ICC (Imaizumi and Hama 1969; Komuro 1989, 1999, 2006; Komuro *et al.* 1999; Yamamoto 1977), and the discovery of their specific marker, the tyrosine kinase receptor (c-Kit), has also contributed to the distinguishing of ICC from other interstitial cells (Maeda *et al.* 1992). ICC are found in the smooth muscle layer of the gut and classified into several subtypes, according to their localization in the muscle layers (Fig. 1). The subtypes are ICC associated with deep muscular plexus of the small intestine (ICC-DMP), myenteric (Auerbach) plexus (ICC-MP), and submuscular plexus of the colon (ICC-SMP), and ICC located within the circular muscle layer (ICC-CM) and within the longitudinal muscle layer (ICC-LM) (Thuneberg 1982, 1989). Physiological investigation of ICC have also shown dramatic progress due to the pacemaker hypothesis proposed by Thuneberg

(Thuneberg 1982, 1989). The major functions of ICC are as pacemakers of gut motility and as intermediates in the transmission of messages between ENS and smooth muscles. Moreover, the function of each subtype of ICC does not seem to be the same depending on the sites where they are localized. It has also become clear that ICC and the ENS almost always work together (Huizinga *et al.* 2011).

However, there have not been any systematic studies on the distribution of ICC subtypes throughout the intestine. Further, many intestinal disorders have been reported to be related to the number or density of ICC. Basic information about ICC, including their actual function in these diseases is not sufficient to be used as diagnostic criteria in each intestinal dysfunction. Therefore, it is very important as basic information, to understand the normal distribution and the features of the normal morphological network in detail, for each subtype of ICC, in different locations of the intestine, in the context of their function in gastrointestinal motility and interaction with ENS.

In the present study, as a systematic study on the distribution of ICC subtypes throughout the intestine, we evaluated the overall distribution pattern of each ICC subtype in the mouse intestine. Based on morphological analyses using specific antibodies to identify ICC (Komuro 1999; Ward *et al.* 1994) and ENS (Fu *et al.* 2013; Iino and Horiguchi 2006), we measured the local density of ICC and compared the difference in the number of ICC between mesenteric and anti-mesenteric sides, and between proximal and distal sites, in the mouse intestine. We also discuss whether the quantitative features of each subtype of ICC may correspond to their possible physiological roles in the mouse intestine.

Materials and Methods

Animals and Ethics

C57BL/6J female mice (6–8 weeks old) were purchased from CLEA Japan (Tokyo, Japan) and used for all experiments designed in this study, which were approved by the Ethical Review Committee of Animal Experiments at Tokyo Women's Medical University (TWMU), approval number is AE18–71 (Mar 31, 2018). Mice were anesthetized using an intraperitoneal injection of ketamine (87 µg/g body weight),




Fig. 1. Subtypes of ICC and their Location. A diagram showing the cross-section of small and large intestine (from submucosal layer to muscle layer) and various subtypes of ICC (in green) and the enteric nervous system (in red). ICC-SMP are present on the inner aspect of the circular muscle layer in the large intestine. ICC-DMP represent a population of cells within the specialized deep muscular plexus region of the small intestine. ICC-CM are located within the circular muscle in the large intestine. ICC-MP are associated with the myenteric (Auerbach) plexus and located between the circular and longitudinal muscular layers throughout the gastrointestinal tract. ICC-LM are located within the longitudinal muscle in the large intestine. ICC-DMP: ICC associated with deep muscular plexus, ICC-SMP: ICC associated with submucosal plexus, ICC-CM: ICC located within the circular muscle layer, ICC-MP: ICC associated with myenteric (Auerbach) plexus, ICC-LM: ICC within the longitudinal muscle layer

xylazine (13 $\mu\text{g/g}$ body weight), and metacam (1 $\mu\text{g/g}$ body weight) and were sacrificed by cervical dislocation. All handling and care of animals was carried out in accordance with the guidelines of the Institute of Laboratory Animals for Animal Experimentation and approved by the Animal Care and Use Committee of TWNU.

Tissue preparation

Tissue samples, including the small intestine, colon, and rectum were removed from at least three mice ($n \geq 3$), placed into phosphate buffered saline (PBS; pH 7.2). The length and the diameter of each intestine was measured and the fecal matter was washed out promptly. For inhibition of contractile activity in the intestines, the intestines were placed into saline solution containing papaverine (0.2 mg/mL), a calcium antagonist, for 10 min at 37°C and then rinsed with PBS. The intestines were inflated with acetone to the original size and fixed in ice-cold acetone for 20 min. After rinsing in PBS, the intestines were opened along the mesenteric border, the posterior side in rectum, and pinned

to a dissecting dish. Lengths of 3 cm of the selected regions (jejunum: the proximal side, ileum: the distal end, colon: proximal colon (1 cm distal from the ileocecal valve), and rectum: the distal end) were prepared and oriented as shown in Fig. 2. In this study, we called the preparations of the jejunum and ileum as the proximal and distal sites of the small intestine, respectively, and the preparations of the colon and rectum as the proximal and distal sites of the large intestine, respectively. The whole-mount preparations of the muscle layers were prepared carefully by peeling off the mucosa and submucosa with fine forceps, under a dissecting microscope.

Immunohistochemistry

Whole-mount samples were placed in PBS containing 0.5% Triton X-100 for 10 min at room temperature (approximately $20 \pm 5^\circ\text{C}$). Non-specific binding was blocked by incubation in 5% Block Ace (DS Pharma Biomedical, Osaka, Japan) in distilled water, for 20 min. The samples were then incubated with a rat monoclonal anti c-Kit antibody (ACK2, OriGene,




Fig. 2. Diagrams of Tissue Preparations. Constant lengths (3 cm) from proximal jejunum, terminal ileum, proximal colon, and rectum were obtained for each preparation and opened along the mesenteric border. (A, B). The whole mount preparations of the muscle layers were made after peeling away the mucosa and submucosa, which were placed in the same direction. The lateral quarter areas were regarded as the mesenteric side and the middle area was regarded as the anti-mesenteric side (C). Photomicrographs were taken in eight random fields from both the mesenteric and anti-mesenteric side of each level of ICC subtype (ICC-DMP, ICC-MP, ICC-SMP, ICC-CM, and ICC-LM) from each part of the intestine (jejunum, ileum, colon, and rectum). The horizontal direction of photomicrographs was in the direction of the long axis.

MD, USA) at a dilution of 1:100, and a polyclonal rabbit UCHL1/PGP9.5 antibody (Proteintech, IL, USA) at a dilution of 1:500, for 2 nights at 4°C, to identify ICC and ENS, respectively. After washing in PBS, the samples were further incubated overnight at 4°C with either Alexa Fluor® 488 conjugated goat anti-rat IgG at a dilution of 1:200 (Eugene, OR, USA), Cy3 conjugated goat anti-rabbit IgG at a dilution of 1:200 (Jackson ImmunoResearch, PA, USA), or Hoechst 33342 at a dilution of 1:2,000 (Dojindo, Kumamoto, Japan) for nuclear staining. All antibodies were diluted in 1% bovine serum albumin in PBS. After washing with PBS, the tissues

were mounted on glass slides and examined with a Zeiss LSM 510 confocal microscope (Zeiss, Germany), at 488 nm (for Alexa Fluor 488), 561 nm (for Cy3), and 405 nm (for Hoechst 33342).

Morphological measurement and statistical analyses

Photomicrographs were taken for eight random fields (400× magnification) from both the mesenteric side and the anti-mesenteric side, at each level of ICC subtype localization (ICC associated with deep muscular plexus (ICC-DMP), ICC

associated with myenteric plexus (ICC-MP), ICC associated with submuscular plexus (ICC-SMP), ICC located within the circular muscle layer (ICC-CM), and ICC located within the longitudinal muscle layer (ICC-LM)), at various sites of the intestine (jejunum, ileum, colon, and rectum), with confocal microscopy. Confocal micrographs were obtained, with digital composites of Z-series, through a depth of 6–12 μm for ICC-DMP, 8–20 μm for ICC-MP, 4–16 μm for ICC-SMP, 4–30 μm for ICC-CM, and 1–5 μm for ICC-LM. Final images were constructed using the Zeiss software (Zen). The number of c-Kit/Hoechst 33342 double labeled cells were counted by examining the confocal micrographs. c-Kit positive areas were measured using ImageJ.

Data were expressed as means (standard deviation: SD) per square millimeter. As an exception, those of ICC-CM and ICC-LM were expressed as means per cubic millimeter because of the thickness of circular and longitudinal muscles in ICC-CM and ICC-LM, respectively, depending on their location. Differences in the data were evaluated by Student's *t*-test. Difference probabilities; $P < 0.05$ was regarded as statistically significant, and $P < 0.01$ was highly significant.

Results

I. Typical morphological features of various subtypes of ICC

The typical features of ICC subtype, as identified by their c-Kit immunoreactivity, at each local site are shown in Fig. 3. In the small intestine, ICC were mainly identified in two regions; the deep muscular plexus (ICC-DMP) and the myenteric plexus (ICC-MP). c-Kit immunoreactivity was rarely found in the thick sublayer of the circular muscle and in the longitudinal muscle layer. ICC-DMP were located adjacent to the nerve bundles and projected long slender processes along the deep muscular plexus (Fig. 3A). In contrast, ICC-MP formed a more complex cellular network in the meshwork of the nerve plexus (Fig. 3B). In the large intestine, four subtypes of ICC were recognized. ICC-SMP were clearly recognized as ICC associated with the submuscular plexus, but in a way, demarcating the submucosal border of the circular muscle layer. Their cell bodies were randomly distributed but connected with each other with their long slender processes (Fig. 3C). In the circular muscle layer, ICC-CM were located within the muscle bundles

(Fig. 3D). They seemed to be closely associated with the nerves within the circular muscle layer that were oriented in parallel with the circular muscle bundles. Their shape was not simply bipolar, but multipolar. However, ICC-MP formed a more complex cellular network (Fig. 3E). Their cell bodies seemed to associate closely to the myenteric plexus and connected with each other with their long slender processes. Within the longitudinal muscle layer, a few ICC-LM were detected as spindle shaped or bipolar cells (Fig. 3F). They ran along the length of the muscle bundles with fewer processes but there were few nerve fiber around.

II. Distribution density of various subtypes of ICC

To investigate the general distribution of ICC in the intestine and to compare the difference in the number of ICC at each site between mesenteric and anti-mesenteric sides, and between proximal and distal sites, we first counted the actual number of ICC at the determined sites in the mouse intestine, as described in Fig. 2.

1) Comparison of distribution density between the mesenteric side and the anti-mesenteric side

In general, no distinctly significant difference in the cell density of most subtypes of ICC, between the mesenteric and the anti-mesenteric side was seen. However, the cell densities of ICC-DMP in the jejunum, ICC-CM in the rectum, and ICC-MP in the rectum were significantly higher at the mesenteric side than the anti-mesenteric side (Table 1).

2) Comparison of distribution density between proximal and distal sites

There was no difference in the cell density of ICC-MP within the small intestine between the proximal and distal sites and that of ICC-SMP and ICC-CM within the large intestine, between the proximal and distal sites. In contrast, ICC-MP in the large intestine showed a significantly higher cell density at the proximal site than in the distal site, and cell density of ICC-DMP in the small intestine and ICC-LM in the large intestine was significantly higher in the distal sites than in the proximal sites (Fig. 4).




Fig. 3. Typical Features of ICC Subtypes in the Mouse Intestine. Whole-mount preparations showing c-Kit positive ICC (green) and PGP9.5 positive nerves (red). Typical features of each ICC subtype are presented as representative subtypes. In the small intestine, ICC-DMP in the ileum (A) and ICC-MP in the jejunum (B), were identified. In contrast, four subtypes of ICC were detected in the large intestine. ICC-SMP in the colon (C) and ICC-CM, ICC-MP, and ICC-LM in the rectum (D-F, respectively). Bar indicates 50 μ m.

Table 1. Mean cell density of ICC in each part of the intestine and the difference between mesenteric and anti-mesenteric sides in the mouse intestine

ICC subtype	part	mesenteric side	anti-mesenteric side	p value
Small intestine				
1. ICC-DMP	jejunum	439 (80.4)	388 (87.5)	0.04 *
(cell/mm ²)	ileum	483 (118.3)	476 (116.3)	0.84
2. ICC-MP	jejunum	721 (129.8)	686 (122.9)	0.34
(cell/mm ²)	ileum	686 (130.3)	664 (124.2)	0.55
Large intestine				
1. ICC-SMP	colon	729 (146.5)	713 (221.4)	0.76
(cell/mm ²)	rectum	719 (116.8)	680 (102.5)	0.22
2. ICC-CM	colon	36411 (15028.3)	29965 (12808.1)	0.11
(cell/mm ³) ^{†1}	rectum	38265 (15378.5)	29302 (14751.3)	0.04 *
3. ICC-MP	colon	1298 (125.1)	1247 (153.7)	0.20
(cell/mm ²)	rectum	960 (169.9)	792 (112.8)	0.0002 **
4. ICC-LM	colon	11457 (7832.7)	12383 (8242.1)	0.69
(cell/mm ³) ^{†1}	rectum	19055 (13326.9)	23416 (19798)	0.37

Data are expressed as mean cell number (standard deviation: SD) per square millimeter.

^{†1} As for exceptions, those of ICC-CM and ICC-LM were expressed as mean cell number per cubic millimeter because of the thicknesses of the circular and longitudinal muscle involving ICC-CM and ICC-LM, respectively, differed by their location. *: $P < 0.05$, **: $P < 0.01$

III. Comparison of the distribution pattern and morphology of ICC-MP between the small intestine and the large intestine

The subtype ICC-MP exists in both the small and large intestine, where it associates with the ENS. Comparison of ICC-MP present in the small intestine and the large intestine was done. The spatial relationship of both types of ICC with the myenteric plexus, can be seen in Fig. 3. The characteristic features between proximal and distal sites are also demonstrated in Fig. 5. Both ICC-MP showed a multipolar shape, with long branching processes that form an interconnecting network. ICC-MP of the small intestine, however, seemed to have thicker branching processes and more lateral branching than ICC-MP of the large intestine.

The cell density of ICC-MP of the large intestine was higher than that of the small intestine, with significant

difference (Table 2). However, the c-Kit positive mean area of ICC-MP per unit area (/1mm²) was larger in the small intestine than the large intestine (Table 2, Fig. 6A). Furthermore, the c-Kit positive mean area of ICC-MP in the proximal site was significantly larger than that in the distal site, for both the small and the large intestine (Table 2, Fig. 6B, C). For comparison, the c-Kit positive area of ICC-SMP did not show any difference between the proximal and distal sites in the large intestine (Table 2, Fig. 6D).

IV. Unique morphological feature of ICC-SMP

Unique feature of ICC-SMP was found in this study. ICC-SMP had flat and dense network. The pattern of network was different between the proximal and distal sites. The connecting pattern of ICC-SMP in the proximal site was mesh-like (Fig. 7A) but that in the distal site was a ladder-




Fig. 4. Mean Area Densities of Subtypes of ICC. The total value for each item from Table 1 were visualized in graphs. The cell number per unit area (1 mm²) were expressed on the Y axis (cells/mm²). * $p < 0.05$, ** $p < 0.01$

like pattern along with a circular muscle layer (Fig. 7B).

Discussion

The aim of this study was to evaluate the distribution pattern of ICC subtypes to understand their role in gastrointestinal motility and interaction with the ENS, depending on their

location. Based on the morphological analyses, using c-Kit and PGP9.5 to identify ICC and ENS, respectively, we measured the normal local density and morphological features of ICC and compared the number of ICC between mesenteric and anti-mesenteric sides, and in between proximal and distal sites using whole mount preparations of the mouse intestine. This approach enabled us to accurately evaluate the distribution density of ICC in the local area.




Fig. 5. Comparison of ICC-MP between the Proximal and Distal sites in the Small and Large Intestine. Photomicrographs show ICC-MP (green) and nerves (red). ICC-MP of small intestine (A, B) and the large intestine (C, D). A: jejunum, B: ileum, C: colon, D: rectum. Bar indicates 50 μ m.

The major and novel findings of this study are presented below.

1) The distribution density of ICC varied depending on the site of the intestine. In general, there was no highly significant difference in cell density of most ICC subtypes between the mesenteric and the anti-mesenteric sides, except for ICC-MP in the rectum. In contrast, highly significant differences, in the distribution density, between proximal and distal sites were seen for ICC-DMP (distal > proximal) in the small intestine and ICC-MP (proximal > distal) and ICC-LM (distal > proximal) in the large intestine, respectively. Notably, the two major pacemakers—ICC-MP in the small intestine and ICC-SMP in the large intestine—showed no difference in cell density between the proximal and distal sites.

2) As a mutually existing subtype of ICC in both small

and large intestines, ICC-MP of the large intestine had a significantly greater distribution density than those of the small intestine, although ICC-MP in the small intestine seemed to have denser network with much thicker processes and more branches than those of the large intestine. The predominance in c-Kit positive area of ICC-MP in the small intestine over those in the large intestine was also seen with a high significance. This indicates their morphologically denser network. Furthermore, in both the small and large intestine, larger c-Kit positive areas were seen in the proximal site than the distal site. In other words, there is a clear gradient of c-Kit positive area of ICC-MP in the intestine, which means that their network, including their processes and branches, is denser in the proximal site than in the distal site.

3) The network pattern of the processes of ICC-SMP in the

Table 2. Difference in cell density and c-kit positive area of ICC-MP between small intestine and large intestine

	ICC subtypes	small intestine		large intestine		<i>p</i> value
Cell Density ^{†1}	ICC-MP (cell/mm ²)	689 (126.4)		1074 (251.3)		<0.0001**
c-kit positive area per unit area (1 mm ²) ^{†2}	ICC-MP (mm ²)	0.374 (0.051)		0.287 (0.057)		<0.0001**
		proximal	distal	proximal	distal	
	ICC-MP (mm ²)	0.384 (0.045)	0.354 (0.052)	0.323 (0.042)	0.245 (0.039)	0.004**
	ICC-SMP (mm ²)			0.270 (0.033)	0.264 (0.028)	0.33

^{†1}: Data were expressed as mean cell number (standard deviation: SD) per square millimeter.

^{†2}: Data were expressed as mean c-kit positive area of ICC (standard deviation: SD) per unit area (mm²). The data of ICC-SMP are also presented for comparison. **: $P < 0.01$




Fig. 6. Comparison of c-Kit Positive Areas (mm²) Per Unit Area (1 mm²). The data in Table 2 were visualized in graphs. c-Kit positive area of ICC-MP per unit area (1 mm²) were expressed on the Y axis (mm²). Note that ICC-MP of the small intestine occupied a larger area than those of the large intestine. The c-Kit positive area of ICC-SMP in the large intestine are also shown for comparison. ** $p < 0.01$, *** $p < 0.0001$




Fig. 7. Morphological Comparison of ICC-SMP between the Proximal and Distal Sites in the Large Intestine. Photomicrographs show ICC (green) and nerve cell elements (red) in the large intestine. ICC-SMP in the proximal site (A) and the distal site (B). ICC-SMP had a flat and dense network. The pattern of network was different in the proximal (colon) and distal sites (rectum). Note that the connecting pattern of proximal ICC-SMP (A) was mesh-like but distal ICC-SMP (B) was ladder-like circumferentially. Bar indicates 50 μ m.

proximal site (colon) of the large intestine was mesh-like, whereas that in the distal site (rectum) was ladder-like. This is the first indication that the ICC-SMP network pattern, between the proximal and distal sites of the large intestine, is different.

The ICC play two major roles in gastrointestinal motility. One is that of a pacemaker and the other is as an intermediate for transmission between the ENS and the smooth muscles. ICC-MP in the small intestine and ICC-SMP in the large intestine have been known to act as pacemaker cells for intestinal motility (Barajas-Lopez and Huizinga 1989; Smith *et al.* 1987; Ward *et al.* 1994). ICC-MP in the small intestine generate pacemaker activity that develops into electrical slow waves (Huizinga *et al.* 1995; Ordog *et al.* 1999; Sanders *et al.* 2014; Ward *et al.* 1994). To the best of our knowledge, the present study is the first to compare the distribution density of ICC-MP between proximal and distal sites, and between mesenteric and anti-mesenteric sides of the murine small intestine. We found that there was no difference in the distribution density of ICC-MP between the proximal and distal sites and between mesenteric and anti-mesenteric sides, with any gradient. The mean distribution density of ICC-MP of small intestine was $689.2 \pm 20.3/\text{mm}^2$ (mean $\pm$ S.E.) in our study, that is close to the result ($725.3 \pm 20.2/\text{mm}^2$) of Mei *et al.* (2009). For over a century, it has been known that the frequency of slow waves and their contractions

decrease from the duodenum to the ileum of the small intestine in rabbits (Alvarez 1914; Maslennikova 1961) and in humans (Christensen *et al.* 1966). The gradient was relative to the distribution density of ENS; therefore, the difference in gastrointestinal motility of a specific region of the intestine was speculated to come from the difference in the density of ENS (Maslennikova 1961). The result of the present study that there is constant distribution density of ICC-MP do not contradict the theory that ENS generates the gradient of contraction frequency in the small intestine. However, the fact that the c-Kit positive area of ICC-MP of the small intestine, was greater at the proximal site than at the distal site, indicates that their denser network in the proximal site could be involved in the gradient of slow wave. In contrast, ICC-SMP is important as a pacemaker that generates slow waves of the large intestine, as reported previously (Barajas-Lopez and Huizinga 1989; Plujà *et al.* 2001; Huizinga *et al.* 2011), and is less influenced by neural inputs (Alberti *et al.* 2005). High frequent contractions are constant all along the colon (Huizinga *et al.* 2011). In this study, we found that the density of ICC-SMP of the large intestine did not have a gradient in the proximal to the distal site and in the mesenteric to the anti-mesenteric side. Previous studies have also reported that ICC-SMP had no gradient in proximal, median, and distal sites of the large intestine in rats (Alberti *et al.* 2007) and mice (Wang *et al.* 2014). Furthermore, for

c-Kit positive area per unit area, there was no difference in the size of ICC-SMP between the proximal and distal sites. We consider that the quantitative feature that ICC-SMP has constant density, support the physiological feature that slow wave of the colon is constant in the full length of large intestine. Since high frequent contractions are less influenced by neural inputs, the density of ICC-SMP might be unrelated to the density of ENS, that is higher at the distal than the proximal site of the large intestine. In contrast, difference in morphological features of ICC-SMP between proximal and distal sites of the large intestine might be important. In the proximal site of the large intestine, the network pattern of the processes of ICC were mesh-like, whereas it was ladder-like in the distal site. However, the functional significance of the different network patterns of ICC-SMP in the large intestine is unclear.

For decades, propagating rhythmic motor patterns in the colon, identical to colonic migrating motor complexes, have been described as either neurogenic (blocked by tetrodotoxin) or myogenic (insensitive to tetrodotoxin) (Huizinga *et al.* 2011). Many studies have indicated the involvement of ICC-MP in the contraction of the large intestine (Alberti *et al.* 2007; Barajas-Lopez and Huizinga 1989; Bayguinov *et al.* 2010; Dickson *et al.* 2010; Liu *et al.* 1998; Plujà *et al.* 2001; Smith *et al.* 1987; Spencer *et al.* 2007). Recently, ICC-MP has been suggested to be the second pacemaker in the large intestine. Huizinga *et al.* (2011) proposed that there are two independent pacemakers in the rat colon, one housed in the ICC-SMP, independent of the ENS, and the other housed in the ICC-MP that is stimulus-dependent, usually dependent on neural activity *in vivo*. Both networks of ICC-SMP and ICC-MP may orchestrate propagating contractions as pacemaker cells in concert with ENS and have a significant role related to ENS. Previous studies have shown that ENS has a greater density in the large intestine than the small intestine. Our results showing a greater distribution density of ICC-MP in the large intestine than the small intestine, as with the density of ENS, is consistent with its important role. However, the fact that ICC-MP in the small intestine seemed to have denser network with much thicker processes and more branches than those of the large intestine might have some relation to the difference in the roles of ICC-MP between the small and large intestines. Further physiological investigation of ICC-MP is needed for the depolarization of the thick

smooth muscle layer in the large intestine.

Smooth muscle cells, ICC, and platelet-derived-growth-factor-receptor-alpha positive cells make up a complex of electrically-coupled cells, known collectively as the SIP syncytium (Sanders *et al.* 2012; Sanders *et al.* 2014). Among all subtypes of ICC, intramuscular ICC (ICC-IM), namely ICC-CM and ICC-LM, in the large intestine are thought to be involved in neuro-effector transduction in GI muscles and have an important role in tonic inhibition. Our results showed that the distribution density of ICC-CM had no gradient between proximal and distal sites but had slight increase at the mesenteric side in the rectum. The average distribution density of ICC-LM was very low as the longitudinal muscle was very thin, but in the distal site (near the anal sphincter), the density of ICC-LM increased with the thickening of the longitudinal muscle. One of the features of ICC-IM was that there were regional dispersions, therefore when we examine the density of ICC-IM, we should check many spots from the preparations. In this study, we found that ICC-LM lie sparsely in the longitudinal muscle and nerve fibers almost never exist in whole mount preparations of the mouse intestine.

Previous studies have indicated that ICC-DMP are innervated by motor neurons and transduce part of the input from enteric motor neurons (Iino *et al.* 2004; Wang *et al.* 2003a; Ward *et al.* 2006). In the present study, there was no gradient of the quantity of nerve bundles on the deep muscular plexus between the proximal and distal sites by PGP9.5 positive area (data not shown). Contrary to the expectation, however, the distribution density of ICC-DMP was higher in the distal site of the small intestine but it did not show the same tendency as the density of ENS. Recently, ICC-DMP has received a great deal of attention in the generation of the segmentation pattern of the small intestine (Huizinga *et al.* 2011). In the distal site of the small intestine, ICC-DMP may facilitate the generation of the rhythmic segmentation motor pattern.

In this study, we found differences in the distribution density between mesenteric and anti-mesenteric sides in ICC-DMP in the jejunum and ICC-CM and ICC-MP in the rectum. Their functional meanings are not clear, although previous studies by Gabella *et al.* have revealed that ENS is distributed at a greater density in the mesenteric side than the anti-mesenteric side. The difference in the distribution of ICC may have some functional relevance to ENS.

Decreasing or abnormal ICC in the large intestine including irritable bowel syndrome (Eshraghian and Eshraghian 2011), Crohn's disease (Wang *et al.* 2007), hypertrophic pyloric stenosis (Huizinga *et al.* 2009), colonic cancer, diverticulitis (He *et al.* 2000), chronic pseudo-obstruction (Struijs *et al.* 2008), and delayed maturation of the ICC in a transient neonatal pseudo-obstruction (Kenny *et al.* 1998) has been demonstrated in various human gastrointestinal motility disorders. Moreover, there exists a discrepancy between reports of decreasing and unchanging of ICC in Hirschsprung disease, known as being caused by abnormal development of the ENS (Horisawa *et al.* 1998; Vanderwinden JM 1996; Yamataka *et al.* 1995). In addition, the possible involvement of other types of cells, such as fibroblasts, macrophages, or mast cells have been suggested for the regulation of smooth muscle movement (Komuro 2012). Therefore, the careful studies on the quantification and distribution of ICC, under normal conditions, are important for the diagnosis and treatment of clinical disorders related to ICC. Although the present study provides fundamental information on the topographical distribution and density of major subtypes of ICC in the mouse intestine, it would lend itself to the consideration of their normal functional and morphological knowledge about ICC in the gastrointestinal tract.

Conclusion

In the present study, we measured the normal local distribution and density of ICC to compare the number of ICC between the mesenteric and anti-mesenteric sides and between the proximal and distal sites, using whole mount preparations of the mouse intestine. The distribution density of ICC varied depending on the site of the intestine. There were gradients of cell density in some subtypes, but the gradient tendency was not always proportional to other subtypes even in the same part of intestine. Namely, there were individual cell densities, morphological features, and network patterns of each subtype depending on the site of the intestine. Therefore, the careful studies on the quantification and distribution of ICC under normal conditions should be important in clinical applications for the diagnosis and treatment of disorders related to abnormal distribution of ICC.

Acknowledgements

The authors are very grateful to Dr. K. Horiguchi, University of Fukui, Faculty of Medical Sciences, for his invaluable encouragement and criticism. The authors would also like to thank Professor Ayako Ishizu, Department of Anatomy, Division of Microscopic Anatomy and Developmental Morphology, for providing the opportunity to complete this study in her laboratory. Special thanks go to Dr. M. Morishima, Dr. S. Kitahara, Dr. K. Shimizu, and Dr. R. Hayakawa for their support and encouragement; Mrs. H. Sagawa, Mrs. K. Motomaru, and Ms. I. Sato for their excellent technical assistance in the Department of Anatomy and Developmental Biology. In this research work we used a Zeiss LSM 510 confocal microscope of Medical Research Institute (MRI), TWNU.

Conflicts of Interest

The authors have no conflict of interest to declare.

References

- Alberti E, Mikkelsen HB, Larsen JO, Jimenez M: Motility patterns and distribution of interstitial cells of Cajal and nitrergic neurons in the proximal, mid- and distal-colon of the rat. *Neurogastroenterol Motil* 17: 133-147 (2005).
- Alberti E, Mikkelsen HB, Wang XY, Diaz M, Larsen JO, Huizinga JD, Jimenez M: Pacemaker activity and inhibitory neurotransmission in the colon of Ws/Ws mutant rats. *Am J Physiol Gastrointest Liver Physiol* 292: 1499-1510 (2007).
- Alvarez WC: Functional variations in contractions of different parts of the small intestine. *Am J Physiol* 35: 177-193 (1914).
- Barajas-Lopez C, Huizinga JD: Different mechanisms of contraction generation in circular muscle of canine colon. *Am J Physiol* 256: G570-580 (1989).
- Bayguinov PO, Hennig GW, Smith TK: Ca²⁺ imaging of activity in ICC-MY during local mucosal reflexes and the colonic migrating motor complex in the murine large intestine. *J Physiol* 588: 4453-4474 (2010).
- Cannon WB: The movements of the intestines studied by means of the Rontgen rays. *J Med Res* 7: 72-75 (1902).
- Christensen J, Schedl HP, Clifton JA: The small intestinal

- basic electrical rhythm (slow wave) frequency gradient in normal men and in patients with variety of diseases. *Gastroenterology* 50: 309-315 (1966).
- Costa M, Furness JB, Gabella G: Catecholamine containing nerve cells in the mammalian myenteric plexus. *Histochemie* 25: 103-106 (1971).
- Daniel EE, Wang YF: Gap junctions in intestinal smooth muscle and interstitial cells of Cajal. *Microsc Res Tech* 47: 309-320 (1999).
- Dickson EJ, Heredia DJ, McCann CJ, Hennig GW, Smith TK: The mechanisms underlying the generation of the colonic migrating motor complex in both wild-type and nNOS knockout mice. *Am J Physiol Gastrointest Liver Physiol* 298: 222-232 (2010).
- Elliott TR, Barclay-Smith E: Antiperistalsis and other muscular activities of the colon. *The Journal of Physiology* 31: 272-304 (1904).
- Eshraghian A, Eshraghian H: Interstitial cells of Cajal: a novel hypothesis for the pathophysiology of irritable bowel syndrome. *Can J Gastroenterol* 25: 277-279 (2011).
- Feldstein AE, Miller SM, El-Youssef M, Rodeberg D, Lindor NM, Burgart LJ, Szurszewski JH, Farrugia G: Chronic intestinal pseudoobstruction associated with altered interstitial cells of cajal networks. *J Pediatr Gastroenterol Nutr* 36: 492-497 (2003).
- Fu YY, Peng SJ, Lin HY, Pasricha PJ, Tang SC: 3-D imaging and illustration of mouse intestinal neurovascular complex. *Am J Physiol Gastrointest Liver Physiol* 304: 1-11 (2013).
- Furness JB, Costa M: Morphology and distribution of intrinsic adrenergic neurones in the proximal colon of the guinea-pig. *Z Zellforsch Mikrosk Anat* 120: 346-363 (1971).
- Furness JB, Costa M: Types of nerves in the enteric nervous system. *Neuroscience* 5: 1-20 (1980).
- Gabella G: Neuron size and number in the myenteric plexus of the newborn and adult rat. *J Anat* 109: 81-95 (1971).
- He CL, Burgart L, Wang L, Pemberton J, Young-Fadok T, Szurszewski J, Farrugia G: Decreased interstitial cell of cajal volume in patients with slow-transit constipation. *Gastroenterology* 118: 14-21 (2000).
- Horisawa M, Watanabe Y, Torihashi S: Distribution of c-Kit immunopositive cells in normal human colon and in Hirschsprung's disease. *J Pediatr Surg* 33: 1209-1214 (1998).
- Huizinga JD, Martz S, Gil V, Wang XY, Jimenez M, Parsons S: Two independent networks of interstitial cells of cajal work cooperatively with the enteric nervous system to create colonic motor patterns. *Front Neurosci* 5: 93 (2011).
- Huizinga JD, Thuneberg L, Kluppel M, Malysz J, Mikkelsen HB, Bernstein A: W/kit gene required for interstitial cells of Cajal and for intestinal pacemaker activity. *Nature* 373: 347-349 (1995).
- Huizinga JD, Zarate N, Farrugia G: Physiology, injury, and recovery of interstitial cells of Cajal: basic and clinical science. *Gastroenterology* 137: 1548-1556 (2009).
- Iino S, Horiguchi K: Interstitial cells of cajal are involved in neurotransmission in the gastrointestinal tract. *Acta Histochem Cytochem* 39: 145-153 (2006).
- Iino S, Ward SM, Sanders KM: Interstitial cells of Cajal are functionally innervated by excitatory motor neurones in the murine intestine. *J Physiol* 556: 521-530 (2004).
- Imaizumi M, Hama K: An electron microscopic study on the interstitial cells of the gizzard in the love-bird (*Uroloncha domestica*). *Z Zellforsch Mikrosk Anat* 97: 351-357 (1969).
- Irwin DA: The anatomy of Auerbach's plexus. *American Journal of Anatomy* 49: 141-166 (1931).
- Kenny SE, Vanderwinden JM, Rintala RJ, Connell MG, Lloyd DA, Vanderhaegen JJ, De Laet MH: Delayed maturation of the interstitial cells of Cajal: a new diagnosis for transient neonatal pseudoobstruction. Report of two cases. *J Pediatr Surg* 33: 94-98 (1998).
- Kitamura Y, Hirota S, Nishida T: Gastrointestinal stromal tumors (GIST): a model for molecule-based diagnosis and treatment of solid tumors. *Cancer Sci* 94: 315-320 (2003).
- Komuro T: Three-dimensional observation of the fibroblast-like cells associated with the rat myenteric plexus, with special reference to the interstitial cells of Cajal. *Cell Tissue Res* 255: 343-351 (1989).
- Komuro T: Comparative morphology of interstitial cells of Cajal: ultrastructural characterization. *Microsc Res Tech* 47: 267-285 (1999).
- Komuro T: Structure and organization of interstitial cells of Cajal in the gastrointestinal tract. *J Physiol* 576: 653-658 (2006).
- Komuro T Atlas of: Interstitial Cells of Cajal in the Gastrointestinal Tract. Springer, Dordrecht, 2012.
- Komuro T, Seki K, Horiguchi K: Ultrastructural characterization of the interstitial cells of Cajal. *Arch Histol Cytol* 62: 295-316 (1999).
- Liu LW, Farraway L, Berezin I, Huizinga JD: Interstitial

- cells of Cajal: mediators of communication between circular and longitudinal muscle layers of canine colon. *Cell Tissue Res* 294: 69-79 (1998).
- Maeda H, Yamagata A, Nishikawa S, Yoshinaga K, Kobayashi S, Nishi K, Nishikawa S: Requirement of c-kit for development of intestinal pacemaker system. *Development* 116: 369-375 (1992).
- Maslennikova LD: [On the relationship between the motor function of the intestines and the gradient of its nervous elements]. *Biull Eksp Biol Med* 52: 117-123 (1961).
- Ordog T, Ward SM, Sanders KM: Interstitial cells of cajal generate electrical slow waves in the murine stomach. *J Physiol* 518: 257-269 (1999).
- Plujà L, Albertí E, Fernández E, Mikkelsen HB, Thuneberg L, Jiménez M: Evidence supporting presence of two pacemakers in rat colon. *Am J Physiol Gastrointest Liver Physiol* 281: 255-266 (2001).
- Ritchie JA: The transport of colonic contents in the irritable colon syndrome. *Gut* 11: 668-672 (1970).
- Saito YA, Stregé PR, Tester DJ, Locke GR, 3rd, Talley NJ, Bernard CE, Rae JL, Makielski JC, Ackerman MJ, Farrugia G: Sodium channel mutation in irritable bowel syndrome: evidence for an ion channelopathy. *Am J Physiol Gastrointest Liver Physiol* 296: 211-218 (2009).
- Sanders KM, Koh SD, Ro S, Ward SM: Regulation of gastrointestinal motility—insights from smooth muscle biology. *Nat Rev Gastroenterol Hepatol* 9: 633-645 (2012).
- Sanders KM, Ward SM, Koh SD: Interstitial cells: regulators of smooth muscle function. *Physiol Rev* 94: 859-907 (2014).
- Smith TK, Reed JB, Sanders KM: Interaction of two electrical pacemakers in muscularis of canine proximal colon. *Am J Physiol* 252: 290-299 (1987).
- Spencer NJ, Bayguinov P, Hennig GW, Park KJ, Lee HT, Sanders KM, Smith TK: Activation of neural circuitry and Ca²⁺ waves in longitudinal and circular muscle during CMMCs and the consequences of rectal aganglionosis in mice. *Am J Physiol Gastrointest Liver Physiol* 292: 546-555 (2007).
- Struijs MC, Diamond IR, Pencharz PB, Chang KT, Viero S, Langer JC, Wales PW: Absence of the interstitial cells of Cajal in a child with chronic pseudoobstruction. *J Pediatr Surg* 43: 25-29 (2008).
- Tana C, Umesaki Y, Imaoka A, Handa T, Kanazawa M, Fukudo S: Altered profiles of intestinal microbiota and organic acids may be the origin of symptoms in irritable bowel syndrome. *Neurogastroenterol Motil* 22: 512-519 (2010).
- Thuneberg L: Interstitial cells of Cajal: intestinal pacemaker cells? *Adv Anat Embryol Cell Biol* 71: 1-130 (1982).
- Thuneberg L: Interstitial cells of Cajal. *Handbook of physiology. The gastrointestinal system, Sect. 6.*, J D Wood American Physiological Society, Bethesda, MD Vol. 1: 349-386 (1989).
- Vanderwinden JM RJ, Liu H, Descamps D: Interstitial cells of Cajal in human colon and in Hirschsprung disease. *Gastroenterology*, pp. 901-910 (1996).
- Wang XY, Chen JH, Li K, Zhu YF, Wright GW, Huizinga JD: Discrepancies between c-Kit positive and Anol positive ICC-SMP in the W/W^v and wild-type mouse colon; relationships with motor patterns and calcium transients. *Neurogastroenterol Motil* 26: 1298-1310 (2014).
- Wang XY, Paterson C, Huizinga JD: Cholinergic and nitrergic innervation of ICC-DMP and ICC-IM in the human small intestine. *Neurogastroenterol Motil* 15: 531-543 (2003a).
- Wang XY, Ward SM, Gerthoffer WT, Sanders KM: PKC-epsilon translocation in enteric neurons and interstitial cells of Cajal in response to muscarinic stimulation. *Am J Physiol Gastrointest Liver Physiol* 285: 593-601 (2003b).
- Wang XY, Zarate N, Soderholm JD, Bourgeois JM, Liu LW, Huizinga JD: Ultrastructural injury to interstitial cells of Cajal and communication with mast cells in Crohn's disease. *Neurogastroenterol Motil* 19: 349-364 (2007).
- Ward SM, Burns AJ, Torihashi S, Sanders KM: Mutation of the proto-oncogene c-kit blocks development of interstitial cells and electrical rhythmicity in murine intestine. *J Physiol* 480 (Pt 1): 91-97 (1994).
- Ward SM, McLaren GJ, Sanders KM: Interstitial cells of Cajal in the deep muscular plexus mediate enteric motor neurotransmission in the mouse small intestine. *J Physiol* 573: 147-159 (2006).
- Yamamoto M: Electron microscopic studies on the innervation of the smooth muscle and the interstitial cell of Cajal in the small intestine of the mouse and bat. *Arch Histol Jpn* 40: 171-201 (1977).
- Yamataka A, Kato Y, Tibboel D, Murata Y, Sueyoshi N, Fujimoto T, Nishiye H, Miyano T: A lack of intestinal pacemaker (c-kit) in aganglionic bowel of patients with Hirschsprung's disease. *J Pediatr Surg* 30: 441-444 (1995).