

An Overview of the Clinical Characteristics of Japanese Patients with Spinal Muscular Atrophy: Data from SMART Consortium

| | |
|-------|---|
| メタデータ | 言語: eng 出版者: 公開日: 2022-02-18 キーワード (Ja): キーワード (En): 作成者: KATO, Tamaki, URANO, Mari, MATSUO, Mari, SAITO, Kayoko メールアドレス: 所属: |
| URL | http://hdl.handle.net/10470/00033123 |

Table S2. Clinical features of 5q SMA.

| type | fasciculation | | respiratory support | | | feeding | scoliosis | |
|---------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|------------------------------|---|
| | tongue | finger | support(-) | NPPV | TPPV | support(+) | scoliosis(+) | surgery |
| | no. of cases (% of cases) | no. of cases (% of scoliosis(+) cases) |
| I (n=105) | 77 (73%) | 12 (11%) | 32 (30%) | 10 (10%) | 63 (60%) | 74 (70%) | 27 (26%) | 1 (4%) |
| II (n=106) | 64 (60%) | 45 (42%) | 67 (63%) | 36 (34%) | 2 (2%) | 5 (5%) | 63 (59%) | 14 (22%) |
| III (n=38) | 4 (11%) | 17 (45%) | 37 (97%) | 1 (3%) | | | 15 (39%) | 1 (7%) |
| total (n=249) | 145 (58%) | 74 (30%) | 136 (55%) | 47 (19%) | 65 (26%) | 79 (32%) | 105 (42%) | 16 (15%) |