

ITGAV polymorphism and disease susceptibility in a Japanese rheumatoid arthritis population

著者名	I IKUNI Noriko, KOBAYASHI Shu, IKARI Katsunori, TOMATSU Taisuke, HARA Masako, YAMANAKA Hisash, KAMATANI Naoyuki, MOMOHARA Shigeki
journal or publication title	Arthritis Research & Therapy
volume	9
page range	405
year	2007-10-31
URL	http://hdl.handle.net/10470/30837

doi: 10.1186/ar2303

Letter

***ITGAV* polymorphism and disease susceptibility in a Japanese rheumatoid arthritis population**

Noriko Iikuni, Shu Kobayashi, Katsunori Ikari, Taisuke Tomatsu, Masako Hara, Hisashi Yamanaka, Naoyuki Kamatani and Shigeki Momohara

Institute of Rheumatology, Tokyo Women's Medical University, 10-22 Kawada, Tokyo, 162-0054 Japan

Corresponding author: Katsunori Ikari, kikari@ior.twmu.ac.jp

Published: 31 October 2007

Arthritis Research & Therapy 2007, **9**:405 (doi:10.1186/ar2303)This article is online at <http://arthritis-research.com/content/9/5/405>

© 2007 BioMed Central Ltd

See related research article by Jacq *et al.*, <http://arthritis-research.com/content/9/4/R63>, and related editorial by Ahnert and Kirsten, <http://arthritis-research.com/content/9/5/108>

A research article by Jacq and colleagues [1], recently published in this journal, reported an association between rheumatoid arthritis (RA) and the rs3738919-C allele in the *ITGAV* gene, which encodes the α_v subunit of the integrin $\alpha_v\beta_3$, in a European Caucasian population.

One feature of the pathophysiology of RA is the hyper-angiogenesis observed in the synovial tissue and, along with excess macrophages and T lymphocytes, $\alpha_v\beta_3$ ligands are also abundant in synovial fluid [2]. As a key player in angiogenesis [3], it has been suggested that *ITGAV* may be a RA susceptibility gene. The linkage study also supports this view because the 2q31 region containing *ITGAV* exhibited linkage in a dense genome scan [4].

Although the European association study by Jacq and colleagues [1] indicated *ITGAV* to be a new minor RA susceptibility gene, a replication study conducted in a population of different ethnicity is helpful in exploring whether the association is caused by a common variance through various ethnic groups. We therefore undertook a large population-based study to investigate the association between *ITGAV* and RA in a Japanese population.

DNA samples were obtained from 1,504 Japanese RA patients and 449 population control individuals [5]. Genotypes were determined using the TaqMan method, in accordance with the manufacturer's instructions (Applied Biosystems, Tokyo, Japan). Table 1 shows the genotype distributions and allelic frequencies of patients and control individuals. Unlike the European population, the rs3738919-C allele was more frequent in control individuals in our Japanese population, and no significant differences were observed in allele frequencies for Japanese RA patients and control individuals (0.908 and 0.922, respectively).

There are several factors that must be accounted for when a study fails to corroborate a previously identified association. One of these is an ethnicity-specific effect. Ethnic differences can result in differences in allele frequencies, and the genetic background of the disease itself may vary between ethnic groups. Another issue that must be considered is that, usually, negative association studies of a target gene outside the HLA region lack sufficient power to detect the genetic effect because the disease-associated genes outside the region may be associated with limited relative risk for disease susceptibility. While considering the odds ratio of Jacq *et*

Table 1**Distribution of *ITGAV* rs3738919 genotypes in Japanese rheumatoid arthritis patients and control individuals**

	Genotype					C versus A		CC + CA vs. AA	
	CC	CA	AA	Total	MAF	OR (95% CI)	<i>P</i> ^a	OR (95% CI)	<i>P</i> ^a
Patients	1,221	259	8	1,488	0.092	1.20 (0.91 to 1.60)	0.23	0.60 (0.16 to 2.72)	0.49
Controls	380	62	4	446	0.078				

The distribution of the genotypes followed the Hardy-Weinberg equilibrium for both groups. ^a*P* values were obtained using Fisher's exact test (R version 2.4.1). CI, confidence interval; MAF, minor allele frequency; OR, odds ratio.

al.'s study [1], our study was designed to have sufficient power to detect the effect of the *ITGAV* gene (>0.96 ; number of cases = 1504, number of controls = 449, OR in the original French study [1] = 1.94, frequency of rs3738919-A allele in the controls = 0.078, significance level = 0.05). However, when the odds ratio is under 1.67, the power is limited to below 0.8, and this may represent a study limitation.

Despite its association with RA in Caucasian populations, an association of the *ITGAV* gene with RA patients in the Japanese was not observed. Further studies in other ethnic groups are necessary to draw definite conclusions.

Competing interests

The authors declare that they have no competing interests.

References

1. Jacq L, Garnier S, Dieude P, Michou L, Pierlot C, Migliorini P, Balsa A, Westhovens R, Barrera P, Alves H, *et al.*: **The *ITGAV* rs3738919-C allele is associated with rheumatoid arthritis in the European Caucasian population: a family-based study.** *Arthritis Res Ther* 2007, **9**:R63.
2. Friedlander M, Brooks PC, Shaffer RW, Kincaid CM, Varner JA, Cheresh DA: **Definition of two angiogenic pathways by distinct alpha v integrins.** *Science* 1995, **270**:1500-1502.
3. Eliceiri BP, Cheresh DA: **The role of alphav integrins during angiogenesis: insights into potential mechanisms of action and clinical development.** *J Clin Invest* 1999, **103**:1227-1230.
4. Osorio YFJ, Bukulmez H, Petit-Teixeira E, Michou L, Pierlot C, Cailleau-Moindrault S, Lemaire I, Lasbleiz S, Alibert O, Quillet P, *et al.*: **Dense genome-wide linkage analysis of rheumatoid arthritis, including covariates.** *Arthritis Rheum* 2004, **50**:2757-2765.
5. Iwamoto T, Ikari K, Inoue E, Toyama Y, Hara M, Yamanaka H, Tomatsu T, Momohara S, Kamatani N: **Failure to confirm association between PDCD1 polymorphisms and rheumatoid arthritis in a Japanese population.** *J Hum Genet* 2007, **52**:557-560.