



LINE-1 Expression is Associated with the Effectiveness of EGFR Inhibitors in Colorectal Cancer Cells

著者名	YAMAMOTO Eiichiro, KAWAKAMI Kazuyuki, HAYASHI Kazuhiko, NAGASHIMA Yoji, NOMURA Minoru, OZAKI Makoto
journal or publication title	Tokyo Women's Medical University Journal
volume	4
page range	52-59
year	2020-12-25
URL	http://hdl.handle.net/10470/00032746

Supplementary Table Relation of histology and stage with location of colorectal cancer.

	n	Location		p-value
		Left	Right	
All cases	98	55	43	
Histology				
tub1	52	26	26	0.21
tub2	33	24	9	
pro	6	3	3	
muc	2	1	1	
NA	5	1	4	
Stage				
I	9	7	2	0.99
II	26	15	11	
III	42	20	22	
IV	17	12	5	
NA	4	1	3	
<i>RAS</i> -wild type	60	37	23	
Histology				
tub1	28	16	12	0.42
tub2	23	17	6	
pro	5	2	3	
muc	2	1	1	
NA	2	1	1	
Stage				
I	8	6	2	0.08
II	17	12	5	
III	25	11	14	
IV	8	7	1	
NA	2	1	1	
<i>RAS</i> -mutant type	38	18	20	
Histology				
tub1	24	10	14	0.20
tub2	10	7	3	
pro	1	1	0	
muc	0	0	0	
NA	3	0	3	
Stage				
I	1	1	0	0.54
II	9	3	6	
III	17	9	8	
IV	9	5	4	
NA	2	0	2	