

LINE-1 Expression is Associated with the Effectiveness of EGFR Inhibitors in Colorectal Cancer Cells

メタデータ	言語: eng 出版者: 公開日: 2021-02-02 キーワード (Ja): キーワード (En): 作成者: YAMAMOTO, Eiichiro, KAWAKAMI, Kazuyuki, HAYASHI, Kazuhiko, NAGASHIMA, Yoji, NOMURA, Minoru, OZAKI, Makoto メールアドレス: 所属:
URL	http://hdl.handle.net/10470/00032746

Supplementary Table Relation of histology and stage with location of colorectal cancer.

	n	Location		p-value
		Left	Right	
All cases	98	55	43	
Histology				
tub1	52	26	26	0.21
tub2	33	24	9	
pro	6	3	3	
muc	2	1	1	
NA	5	1	4	
Stage				
I	9	7	2	0.99
II	26	15	11	
III	42	20	22	
IV	17	12	5	
NA	4	1	3	
<i>RAS</i> -wild type	60	37	23	
Histology				
tub1	28	16	12	0.42
tub2	23	17	6	
pro	5	2	3	
muc	2	1	1	
NA	2	1	1	
Stage				
I	8	6	2	0.08
II	17	12	5	
III	25	11	14	
IV	8	7	1	
NA	2	1	1	
<i>RAS</i> -mutant type	38	18	20	
Histology				
tub1	24	10	14	0.20
tub2	10	7	3	
pro	1	1	0	
muc	0	0	0	
NA	3	0	3	
Stage				
I	1	1	0	0.54
II	9	3	6	
III	17	9	8	
IV	9	5	4	
NA	2	0	2	