

Enteral administration of antiepileptic agents could have efficacy for prevention of post-traumatic seizures in severe traumatic brain injury

メタデータ	言語: eng 出版者: 公開日: 2021-10-19 キーワード (Ja): キーワード (En): 作成者: MOCJIDUKI, K , SUZUKI, H , KAWASAKI, A , HOTTA, M , NAMIKI, M , HARADA, T , TAKEDA, M , YAGUCHI, A メールアドレス: 所属:
URL	http://hdl.handle.net/10470/00032989

This work is licensed under a Creative Commons Attribution 3.0 International License.






Figure 1 (abstract P479). Calibration.

Results We studied 531 patients, age 40.35 ± 19.75 years, APACHE II 17.94 ± 6.97 points, Glasgow Coma Scale at admission 7.53 ± 3.83 points. Cranial tomography at admission was: diffuse injury type I (10.4%), diffuse injury type II (28.1%), diffuse injury type III (24.5%), diffuse injury type IV (8.3%), evacuated mass lesion (22.6%), non-evacuated mass lesion (6.2%). Hospital mortality was 28.6%, 171 (32.2%) patients died after 1 year (6.6% missing) and 181 (34.1%) died after 3 years




Figure 2 (abstract P479). Discrimination.

(16.2% missing). Regarding work activities, after 1 year, 28.5% of 326 patients evaluated have no difficulties with work, 4.6% have difficulties but work as before, 10.1% work only part-time or have changed to a job requiring minimum effort and 56.7% of patients do not work. After 3 years, 41.2% of 238 patients evaluated have no difficulties with work, 4.6% have difficulties but work as before, 12.6% work only part-time or have changed to a job requiring minimum effort and 41.6% of patients do not work. Evolution between 1 and 3 years by the McNemar test was statistically significant ($P < 0.001$). A total of 173 patients were in similar situation, only one had deteriorated and in 62 (26.05%) patients the evaluation of work activity had improved.

Conclusion After 1 year of admission from the ICU with TBI, more than 50% of patients have difficulties with work. After 3 years the number of patients who work has increased although approximately 40% of the surviving patients do not work.

P481

Enteral administration of antiepileptic agents could have efficacy for prevention of post-traumatic seizures in severe traumatic brain injury

K Mocjiduki, H Suzuki, A Kawasaki, M Hotta, M Namiki, T Harada, M Takeda, A Yaguchi

Tokyo Women's Medical University, Tokyo, Japan

Critical Care 2014, 18(Suppl 1):P481 (doi: 10.1186/cc13671)

Introduction Antiepileptic prophylaxis is recommended for preventing only early post-traumatic seizures (PTS) in the guidelines for the management of severe traumatic brain injury (TBI) by the Brain Trauma Foundation. Phenytoin is recommended to reduce the incidence of early PTS prophylaxis. Early enteral nutrition has recently shown theoretical advantages for prevention of bacterial translocation to maintain normal turnover of gut mucosa and is commonly used for TBI patients. Our hypothesis is that the enteral administration of antiepileptic agents is also useful for early PTS.

Methods This retrospective observational study included all adult patients admitted to our tertiary academic medico-surgical ICU due to TBI from September 2011 to August 2012. Patients who have epilepsy as a past history were excluded. Clinical data were collected from electrical medical archives. The baseline characteristics collected were age, gender, diagnosis, antiepileptic agents, timing of start and adverse effects of those agents, and methods of administration.

Results Of 65 patients with TBI, 25 patients (18 men, seven women; mean age 56.7 ± 20.1) who were administered antiepileptic agents for PTS prophylaxis were studied. Fifteen cerebral contusions, 10 acute subdural hematomas, nine traumatic subarachnoid hemorrhages, two cerebral infarctions, two pneumocephalus and one traumatic intracerebral hemorrhage were shown in 25 patients. All patients were alive 28 days after the injury. Fourteen patients (56%) were intravenously administered (13 phenytoin and one phenobarbital), while 11 patients (44%) were administered with enteral feeding (four valproates, four carbamazepine and three zonisamides) as PTS prophylaxis. The average start day of PTS prophylaxis was 2.6 days after the injury by intravenous administration, and 2.2 days by enteral administration, respectively. Two patients with phenytoin showed hepatic dysfunction as an adverse effect and no patient showed early PTS both by intravenous and by enteral administrations.

Conclusion The present study has some limitations because it is a single-center retrospective analysis. However, enteral administration of antiepileptic agents could be useful for PTS prophylaxis. Considering cost, adverse effects and serum monitoring, there is a possibility of enteral administration of antiepileptic agents as an alternative to intravenous phenytoin.

P482

Simulation-based education for cardiopulmonary resuscitation and airway management protocols: a brief report of a systematic review and meta-analysis

A Cortegiani, V Rusotto, A Naro, G Sanzo, G Grutta, SM Raineri, A Giarratano
University of Palermo, Italy
Critical Care 2014, 18(Suppl 1):P482 (doi: 10.1186/cc13672)

Introduction We aimed to summarize the efficacy of simulation-based education in cardiopulmonary resuscitation and airway management [1].

Methods We searched the MEDLINE, Scopus and EMBASE databases for all peer-reviewed articles enrolling physicians/medical students in a simulation of either cardiopulmonary resuscitation or airway management protocols compared with no intervention or traditional teaching methods. We categorized the outcomes of the studies into four groups: task success, process skill, time skill, knowledge. Task success was defined as evaluation of successful completion of the task, process skill as evaluation of the procedure, time skill as the time required to complete the task, and knowledge as the objective assessment of conceptual understanding. When studies investigated more than one outcome, we considered the primary outcome, the overall measure or the most clinically relevant outcome.

Results From 8,528 articles, we selected 24 studies (13 randomized controlled studies, eight pre-post studies, three case-control studies) involving 1,149 participants. Compared with no intervention or traditional teaching methods, simulation was associated with a significant improvement from mild to moderate of all outcomes (Figure 1). Log of odds ratio for task success was 2.03 (0.46 to 3.59) in

favor of simulation. Pooled effect size for process skill was 0.48 (0.11 to 0.84), for time skill was 0.29 (0.13 to 0.73) and for knowledge was 0.41 (0 to 0.84).

Conclusion Simulation is an effective educational method to improve performance of physicians/medical students in the application of protocols for cardiopulmonary resuscitation and airway management.

Reference

1. McGaghie WC, et al.: A critical review of simulation-based medical education research: 2003–2009. *Med Educ* 2010, 44:50-63.

P483

Video analysis of cardiopulmonary resuscitation performance of ambulance crews during transportation

H Giga¹, T Otani¹, T Sadamori¹, K Une¹, Y Kida¹, K Ota¹, J Itai¹, S Yamaga¹, S Kusunoki², S Ohshimo¹, Y Iwasaki¹, N Hirohashi¹, K Tanigawa¹
¹Hiroshima University, Hiroshima, Japan; ²Critical Care Medical Center, Hiroshima Prefectural Hospital, Hiroshima, Japan
Critical Care 2014, 18(Suppl 1):P483 (doi: 10.1186/cc13673)

Introduction High quality of chest compressions during cardiopulmonary resuscitation (CPR) is a critical determinant of outcome from out-of-hospital cardiac arrest (OHCA). Unfortunately, however, victims often do not receive adequate chest compression for various reasons, particularly during transportation. Recent studies have demonstrated the interruption time of chest compression using transthoracic impedance analysis, but more information is needed to evaluate the performance of CPR provided by ambulance crews and reveal reasons for hands off chest during CPR.

Methods All ambulances of the Hiroshima City Fire Department are equipped with a specially designed transmission device (RVT-SD200; Sony) that transmits high-resolution visual images and patient vital data using video cameras and a bio-monitor. We analyzed video data of OHCA patients transported by ambulance from November 2012 through December 2012, and evaluated the performance of CPR during transportation in accordance with the 2010 guidelines. The hands-off time was calculated as the time without chest compressions divided by the total CPR time.

Results Thirty-two resuscitation episodes during transportation by ambulance were analyzed. Median CPR time per episode was 846 seconds (range 126 to 1,833 seconds). In total, the fraction of time without chest compression was $19.5 \pm 7.6\%$ (mean \pm SD). Reasons for interruption and its fraction of time in total hands-off time were as follows: 36% accounted for rhythm analysis/pulse check, 31% for ventilation, 11% for setting up automated chest compression devices, 8% for tracheal intubation/placement of supraglottic airway devices, 4% for intravenous line placement/administration of adrenaline, 3% for rescuer change, and 7% for adjustment of patient position/correction of rescuer posture and others.

Conclusion The fraction of time without chest compression observed in this study was comparative with those found in other studies in spite of the difficult situations, such as during transportation. Most frequent reasons for hands-off time were rhythm analysis and ventilation even though the ambulance crews strictly adhered to the guidelines.

